



Determinants of Energy Intake from Ultra-Processed Food Consumption among Adolescent Girls in Tasikmalaya City, Indonesia

Lilik Hidayanti^{1*}, Dian Saraswati² and Rizka Fikrinnisa¹

¹Nutrition Study Program, Faculty of Health Science, Universitas Siliwangi, Tasikmalaya, Indonesia; ²Public Health Study Program, Faculty of Health Science, Universitas Siliwangi, Tasikmalaya, Indonesia

Received: November 15, 2025; **Revised:** February 20, 2026; **Accepted:** February 20, 2026

Abstract

Adolescence is a critical period characterized by significant physical, cognitive, psychosocial, and emotional changes that influence dietary behaviors, including increased consumption of ultra-processed foods (UPFs), which may contribute to chronic disease risk. This study aimed to analyze the determinants of energy intake from UPFs among adolescent girls in Tasikmalaya City. This observational study employed a cross-sectional design, involving 514 adolescent girls from 10 public junior high schools in Tasikmalaya, selected through proportional random sampling. The energy intake from UPFs was collected using 2×24-hour food recalls, then separated using Nova 4. Information on potential determinants, including individual and interpersonal factors, was obtained through a structured questionnaire. Data were analyzed using the chi-square test and logistic regression. The results showed that one-fifth (21%) of participants had energy intake dominated by UPFs. Significant individual factors associated with the dominance of energy intake from UPFs included screen time duration ($p = 0.001$; OR = 2.424 [95% CI = 1.420-4.140]) and main meals ($p = 0.019$; OR = 1.996 [95% CI = 1.121-3.554]). Interpersonal factors related to the dominance of energy intake from UPFs were family status ($p = 0.006$; OR = 2.336 [95% CI = 1.275-4.282]). Screen time duration emerged as the strongest contributing factor. In conclusion, dominant energy intake from UPFs among adolescent girls is influenced by screen time duration, main meals, and family status. Parents need to pay attention to adolescent girls' nutritional intake, especially limiting UPFs consumption. Policies regarding healthy snacks sold in school cafeterias also need to be implemented.

Keywords: adolescent girls; dietary patterns; energy intake; screen time; ultra-processed foods

INTRODUCTION

Adolescence is a phase of life characterized by significant changes in physical, cognitive, psychosocial, and emotional development, as well as health problems distinct from those of childhood and adulthood (Singh et al., 2019). Worldwide, in 2019, there were 1.2 billion teenagers aged 10 to 19 years, and 90% of them lived in low- and middle-income countries (LMICs) (United Nations, 2019). Therefore, improving the health and well-being of

adolescents, especially adolescent girls in developing countries, is a global health priority (Sparrow et al., 2021; Shinde et al., 2023). Unfortunately, adolescent girls in developing countries tend to have poor eating habits (UNICEF Indonesia, 2021).

The habit of eating outside is common among young people (Wellard-Cole et al., 2021). The high frequency of eating outside increases energy, fat, sugar, and sodium intake (Putri and Anwar,

* **Corresponding author:** lilikhidayanti@unsil.ac.id

Cite this as: Hidayanti, L., Saraswati, D., & Fikrinnisa, R. (2026). Determinants of Energy Intake from Ultra-Processed Food Consumption among Adolescent Girls in Tasikmalaya City, Indonesia. *AgriHealth: Journal of Agri-food, Nutrition and Public Health*, 7(1), 44-58. doi: <http://dx.doi.org/10.20961/agrihealth.v7i1.111066>

2024), as the food usually consumed outside is ultra-processed foods (UPFs) (Andrade et al., 2020; Souza et al., 2022). Consumption of UPFs contributes to 50% of total energy intake in some Northern European countries and 25% in some developing countries (Sandoval-Insausti et al., 2020). From 2007 to 2022, there was an increase in UPFs consumption, marked by higher sales in South Asian, Southeast Asian, and Sub-Saharan African countries (Monteiro et al., 2025). Indonesia, one of the developing countries in Southeast Asia, is also experiencing an increase in UPFs consumption (Colozza et al., 2023), with one-third of the Indonesian population exceeding the recommended intake of sugar, salt, and fat (Dewanti et al., 2022).

UPFs are typically high in sugar, salt, and fat (Monteiro et al., 2025) and are major contributors to adolescents' energy intake (Muraro et al., 2023). Young people (ages 12 to 19) consume more energy from UPFs than children (ages 2 to 11) (CDC, 2020). The World Health Organisation (WHO) stated in 2020 that 80% of young people worldwide often consume UPFs, including 50% at lunch, 15% at dinner, and 15% at breakfast (Yetmi et al., 2021). Increased UPFs consumption correlated with increased intake of free sugars, total fat, and saturated fat, and decreased fiber, protein, potassium, zinc, and magnesium, as well as vitamins A, C, D, E, B12, and niacin (Martini et al., 2021).

High UPFs consumption is associated with a higher risk of health deterioration, especially cardiometabolic disease, increased risk of mental disorders, and increased risk of death (Lane et al., 2024). Other findings suggest that high UPFs consumption is associated with a higher risk of diabetes, hypertension, dyslipidemia, and obesity (Vitale et al., 2024). According to a national health survey, the prevalence of non-communicable diseases in Indonesia has been increasing annually (Ministry of Health of Indonesia, 2023). Food consumption, suspected to be a risk factor for non-communicable diseases, includes increased intake of sugar, salt, and excess fat (Andarwulan et al., 2021; Thapsuwan et al., 2024), which are widely present in UPFs (Kesaite et al., 2024). Efforts to control UPFs consumption among adolescent girls can be achieved by identifying the determinants of high UPFs intake.

The social ecological model (SEM) approach, which includes individual, interpersonal, and

environmental factors, is a valuable conceptual framework for understanding health behavior (McLeroy et al., 1988). Individual factors include knowledge, personal behavior and consumption; interpersonal factors include the influence of the social environment, such as family; and environmental factors include community and policy. The SEM approach has been widely used to describe the determinants of unhealthy food consumption practices, such as high-salt intake (Mensah et al., 2025). In UPF's consumption behavior, individual and interpersonal factors are essential drivers (Vashtianada et al., 2023), and these factors influence choice (Santos and Assunção, 2025).

This research was conducted in Tasikmalaya, an area in West Java Province that has the largest population in Indonesia (Provincial Government of West Java, 2024). The population of Tasikmalaya is dominated by the Sundanese ethnic group, known for their consumption of plant-based foods (Raihandhany and Purnomo, 2025). However, research shows that consumption of UPFs is increasing globally, not only in urban areas but also in rural areas (Choudhury et al., 2025). This study aims to analyze the determinants of UPFs using the SEM approach, with a focus on individual and interpersonal factors in Tasikmalaya City, an area with food consumption dominated by plant-based foods.

MATERIALS AND METHOD

Study design and participants

This study employed an observational, cross-sectional design. The population in this study was female junior high school students in Tasikmalaya City, Indonesia. Tasikmalaya City has 21 public junior high schools distributed across 10 sub-districts. Ten public junior high schools were selected as research sites, one from each sub-district. In Tasikmalaya City, there are 6 sub-districts with more than 1 public junior high school, so the sample will be selected randomly using a random picker. The calculation of the minimum sample size, with a 25% non-response rate, yielded 535 participants. The minimum sample size is calculated using Equation 1 (Mukti, 2025).

$$n = \frac{N}{1 + Ne^2} \quad (1)$$

Where, n = Sample, N = Population, and e = Error (10%).

Participants were selected using proportional random sampling. At the end of the study, 21 respondents did not participate in the second food recall and were excluded, leaving 514 participants who could be analyzed. Before the study was conducted, the participants and their parents or legal guardians signed an informed consent form.

Data collection

Fifteen eighth-semester nutrition students participated in data collection for this study. The collected data included both dependent and independent variables. The dependent variable was energy intake from UPFs, obtained by comparing energy intake from UPFs with that from non-UPFs. The results were grouped into 2 categories: UPFs dominance and non-UPFs dominance. UPFs dominance occurred when energy from UPFs was greater than that from non-UPFs. UPF intake was obtained by subtracting foods included in the UPF group from total daily food intake. UPFs were industrial formulations made from inexpensive ingredients extracted from or derived from whole foods, combined with additives. Both local and commercial products were determined using the Nova 4 classification (Monteiro et al., 2019). Total daily food consumption was obtained from a non-consecutive 2×24-hour food recall (Huang et al., 2022) conducted on weekdays and holidays, using a food photo book to help participants estimate household size.

The independent variables were determinants of UPFs, based on a social ecological approach, including individual, interpersonal, environmental, and policy factors. Environmental and policy factors were assumed to be homogeneous because the research was conducted in public junior high schools within the same area (Tasikmalaya City). Therefore, the determinants of energy intake from UPFs in this study included individual factors and interpersonal factors. Individual factors included age, grade, pocket money, sleep duration, late-night sleep, screen time duration, physical activity, nutritional literacy, main eating habits, and breakfast habits. Meanwhile, interpersonal factors included parents' education and occupation, the number of family members, and family status.

Age is the time span in years from the participant's birth until the research was conducted, grouped into 12 to 13 and 14 to 15. Grade is the educational level at which the research was conducted, divided into VII, VIII, and IX. Pocket money is the participant's answer to the question of how much money (IDR) they receive in one day, divided into $> 20,000$ and $\leq 20,000$ IDR per day (Hidayanti et al., 2023). Sleep duration is the time span in hours from when the participants went to sleep until they woke up yesterday, divided into < 8 and ≥ 8 hours (Cortes et al., 2023; Grasaas et al., 2024). Late-night sleeping is the time at which the participant begins to fall asleep at night, grouped as "yes" if the participant went to sleep ≥ 11.00 PM and "no" if they went to sleep < 11.00 PM (Zhong et al., 2025). Screen time duration is the length of time (hours) the participant spends in front of the gadget in one day, divided into < 4 and ≥ 4 hours (Zablotsky et al., 2025).

Physical activity is the participant's answer to the question of how many times they exercise in one week, with a minimum duration of 30 minutes, divided into < 3 and ≥ 3 times per week (James et al., 2023). Nutritional literacy is the participant's answer score on questions based on the Diet and Health Knowledge Survey (DHKS) questionnaire (Bahramfard et al., 2020), grouped into "bad" ($< 50\%$ correct answers) and "good" ($\geq 50\%$ correct answers). The main meals are the frequency (times per day) of consuming complete meals consisting of staple foods and side dishes at the main times of the day (morning, noon, and evening) (Ishikawa-Takata et al., 2021). Main meals were categorized into < 3 and ≥ 3 times per day. Breakfast habits are the frequency (times per week) of food consumption from 05.00 AM to 09.00 AM, which was categorized as "frequent" (3 times per week) or "rare" (< 3 times per week) (Gutkowska et al., 2025).

Parental education is the formal education completed by the father and mother, divided into elementary school, junior high school, senior high school, and university. Parental occupation refers to the activities carried out by parents to earn money, divided into working and non-working activities. The number of family members is the number of people living in the same household as the participant, grouped into > 4 and ≤ 4 (Laksono and Wulandari, 2021). Family status is the

condition of the participant's family at the time the research was conducted, grouped as "complete" if the participant lives with a complete family (father and mother), and "incomplete" if the participant lives only with the father or mother, or neither.

Statistical analysis

Data analysis included univariate, bivariate, and multivariate analyses. In the univariate analysis, data were presented in a frequency distribution table, along with mean, SD, minimum, and maximum values. The relationship between determinant factors and energy intake from UPFs was tested using the chi-square test, with significance set at $p < 0.05$. To fulfil the 2×2 table in the chi-square test, variables with more than 2 categories were changed into 2 categories. The variable of parental education level was grouped into 2 categories based on the 9-year compulsory education program (elementary-junior high school and senior high school-university) (Republic of Indonesia, 2003). The grouping of grade variables was changed to grades VII and VIII-IX. In the bivariate analysis, variables with $p < 0.25$ were further tested using logistic regression modelling.

Ethical consideration

This study has received ethical approval under No. 004616/UNIVERSITAS DIAN NUSWANTORO/2025, dated September 1, 2025.

RESULTS AND DISCUSSION

The total daily nutritional intake of the participants showed that the mean total energy intake was $1,363.3 \pm 459.3$ kcal, total fat intake was 49.2 ± 25.5 g, total carbohydrate intake was 189.6 ± 90.08 g, and total sodium intake was $1,052.2 \pm 737.6$ mg (Table 1). For UPFs consumption, the mean energy, fat, and carbohydrate intake derived from UPFs was lower than from non-UPFs (493.4 ± 286.8 vs 869.3 ± 375.7 kcal; 17.8 ± 12.4 vs 30.6 ± 16.2 g; and 67.0 ± 42.6 vs 122.5 ± 81.5 g, respectively). In contrast, the mean sodium consumption from UPFs (829.2 ± 650.9 mg) was higher than from non-UPFs (222.9 ± 368.2 mg) (Table 1). Of the 514 participants, one-fifth had energy (21.6%), fat (25.7%), and carbohydrate (23.0%) intake from UPFs that dominated daily intake. Meanwhile, more than two-thirds of the participants (84.2%) obtained the majority of their daily sodium intake

from UPFs (Table 1). The use of a 2×24 -hour recall may not fully reflect daily food intake, but this research attempted to increase data validity by conducting the recall on non-consecutive school days and holidays. Potential recall bias and challenges in estimating portion sizes were addressed by using a food photo booklet to assist participants in estimating household measures.

Globally, UPFs consumption is increasing, especially among children and adolescents. This increase occurs not only in high-income countries but also in middle-income countries, such as Indonesia (Mescoloto et al., 2024). In this study, one-fifth of participants (21.6%) consumed more energy from UPFs than non-UPFs sources. Consumption of UPFs contributes to significant increases in energy (Shim et al., 2022), fat (Costa et al., 2024), carbohydrates (Bestari et al., 2023), and sodium (Cortes et al., 2023). Therefore, higher consumption of UPFs also increases the risk of chronic diseases (Matos et al., 2021), such as coronary heart disease (Li et al., 2023), diabetes mellitus (Moradi et al., 2021), and hypertension (Wang et al., 2022).

In this study, participants' sodium intake was primarily from UPFs. UPFs have become a popular snack in developing countries (Moodie et al., 2021), including Indonesia (Nurhasan et al., 2024). Rapidly, high consumption of UPF has become a public health concern due to the high sodium content (Islam et al., 2022). The average salt consumption among Indonesians is 6.6 g, equivalent to 2,674 mg of sodium per day, and the highest sodium content is in snack chips ($1,394.4$ mg per 100 g) (Pradana et al., 2025). In Tasikmalaya, there are local spices such as *atom bulan*, *atom bintang*, *aida*, and a unique onion sauce that has a salty taste (Zahra, 2023). These spices are often added to schoolchildren's snacks, such as sausages, nuggets, fried macaroni, *seblak*, *basreng*, *baso aci*, etc.

Table 2 presents individual and interpersonal factors of adolescent girls in Tasikmalaya City. Regarding individual factors, the mean age of participants was 13 ± 0.8 years, and most were in eighth grade. The majority (81.3%) had pocket money of $\leq 20,000$ IDR per day. More than half (51.0%) had nutritional literacy levels in the bad category. Furthermore, more than half of the participants had a sleep duration of < 8 hours per day (59.5%) and started sleeping after 11.00 PM (12.1%). In addition, most participants spent

Table 1. Nutrient intake from UPF in adolescent girls in Tasikmalaya City

Nutrition content in UPF	n (%)	Mean±SD	Min-max
Energy (kcal)			
Energy total		1,363.3±459.3	364.0-3,976.7
Energy from UPF		493.4±286.8	0-2,162.2
Energy from non-UPF		869.3±375.7	0-2,359.4
Energy domination			
UPF	111 (21.6)		
Non-UPF	403 (78.4)		
Fat (g)			
Fat total		49.2±25.5	5.8-402.4
Fat from UPF		17.8±12.4	0-117.3
Fat from non-UPF		30.6±16.2	0-113.0
Fat domination			
UPF	132 (25.7)		
Non-UPF	382 (74.3)		
Carbohydrates (g)			
Carbohydrates total		189.6±90.08	48.1-1,444.1
Carbohydrates from UPF		67.0±42.6	0-326.5
Carbohydrates from non-UPF		122.5±81.5	0-1,348.7
Carbohydrates domination			
UPF	118 (23.0)		
Non-UPF	396 (77.0)		
Sodium (mg)			
Sodium total		1,052.2±737.6	12.9-7,043.6
Sodium from UPF		829.2±650.9	0-3,828.6
Sodium from non-UPF		222.9±368.2	0-6,442.1
Sodium domination			
UPF	433 (84.2)		
Non-UPF	81 (15.8)		

> 4 hours per day on screens (68.9%) and engaged in physical activity or exercise < 3 times per week (84.8%). In terms of eating habits, most participants consumed main meals < 3 times per day (52.1%) and rarely (73.7%) ate breakfast (Table 2).

In interpersonal factors, the majority of fathers (47.9%) and mothers (47.1%) had a senior high school education, the majority of fathers (96.1%) were employed, and the majority of mothers (73.9%) were unemployed. The average number of family members was 4±1, with the majority (53.3%) having fewer than 4 members, and 88.9% having complete families (father and mother still together) (Table 2).

The result of statistical analysis in Table 3 showed that the individual factors, including age, grade, pocket money, sleep duration, late-night sleep, physical activity, and nutritional literacy, were not related to the dominance of energy

intake from UPFs. Meanwhile, the screen time, main meals, and breakfast habits were associated with the dominance of energy intake from UPFs ($p = 0.001$, 0.005 , and 0.003 , respectively).

In the interpersonal factors, variables for parents' education and occupation, and family members were not related to the dominance of energy intake from UPFs ($p > 0.05$). Only family status showed a significant relationship with the dominance of energy intake from UPFs ($p = 0.035$; OR = 1.984; 95% CI = 1.093-3.601).

At the beginning of the modelling, 7 variables met the inclusion criteria ($p < 0.25$). These consisted of 4 individual factors (screen time duration, nutritional literacy, main meals, and breakfast habits) and 3 interpersonal factors (maternal education, maternal occupation, and family status). Furthermore, logistic regression modelling was carried out by removing, one by one, variables with $p > 0.05$, starting with the

Table 2. Individual and interpersonal factors of adolescent girls in Tasikmalaya City

Variable	n (%)	mean±SD	Min	Max
Individual factors				
Age (years)		13±0.8	12	15
12-13	227 (44.2)			
14-15	287 (55.8)			
Grade				
Complete category				
VII	179 (34.8)			
VIII	217 (42.2)			
IX	118 (23.0)			
Two categories				
VII	179 (34.8)			
VIII-IX	335 (65.2)			
Pocket money (IDR per day)		18,800±7,700	2,000	50,000
> 20,000	96 (18.7)			
≤ 20,000	418 (81.3)			
Sleep duration (hours per day)		7±1	3	9.5
< 8	306 (59.5)			
≥ 8	208 (40.5)			
Late-night sleeping				
Yes	62 (12.1)			
No	452 (87.9)			
Screen time duration (hours per day)		7.6±4.3	2	18
> 4	354 (68.9)			
≤ 4	160 (31.1)			
Physical activity (times per week)		1.7±1.2	0	6
< 3	436 (84.8)			
≥ 3	78 (15.2)			
Nutrition literacy		13.2±3.4	3	22
Bad	262 (51.0)			
Good	252 (49.0)			
Main meals (times per day)		2±1	0	5
< 3	379 (73.7)			
≥ 3	135 (26.3)			
Breakfast habits (times per week)		3±2	0	7
Rare	268 (52.1)			
Frequent	246 (47.9)			
Interpersonal factors				
Father's education				
Complete category				
Elementary	101 (19.6)			
Junior high school	88 (17.1)			
Senior high school	246 (47.9)			
University	79 (15.3)			
Two categories				
Elementary-Junior high school	189 (36.8)			
Senior high school-university	325 (63.2)			
Mother's education				
Complete category				
Elementary	99 (19.3)			

Table 2. *Continued*

Variable	n (%)	mean±SD	Min	Max
Junior high school	101 (19.6)			
Senior high school	242 (47.1)			
University	72 (14.0)			
Two categories				
Elementary-Junior high school	200 (38.9)			
Senior high school-university	314 (61.1)			
Father's occupation				
Not working	20 (3.9)			
Working	494 (96.1)			
Mother's occupation				
Working	134 (26.1)			
Not working	380 (73.9)			
Family members (persons)		4±1	2	9
> 4	235 (47.5)			
≤ 4	279 (54.3)			
Family status				
Incomplete	57 (11.1)			
Complete	457 (88.9)			

variable that had the most significant *p*-value (mother's education). When the mother's education, occupation, and nutritional literacy variables were removed, none of the OR values changed by > 10%. However, when the breakfast habit variable was removed, the ORs changed by > 10% for all variables. Therefore, the variables of maternal education, maternal occupation, and nutritional literacy remain excluded from the model, while breakfast habits were re-entered. In the final model, 3 variables (screen time duration, main meals, and family status) were associated with the dominance of energy intake from UPFs, and screen time duration was the most influential factor (Table 4). However, this study has not considered the influence of peer groups and the presence of street food around schools.

Consumption of UPFs is increasing among adolescents across various countries and demographics (Gonçalves et al., 2023). In this study, participants who spent ≥ 4 hours per day in front of screens had higher energy consumption from UPFs than those who did not, with statistical tests indicating a relationship between screen time duration and energy intake from UPFs. Research involving Brazilian adolescents also showed that adolescents who spent more time in front of screens had a higher percentage of total energy intake from UPFs (Rocha et al., 2021).

Adolescents' screen time is primarily spent accessing social media for various purposes,

including maintaining social connections, seeking information, and expressing themselves (Ouvrein et al., 2019). The majority of adolescents use social media daily, and many report being online almost constantly (Pereira et al., 2024). While using various social media applications, they are also likely to be exposed to UPF's advertising. This is because UPFs advertising is on the rise globally, especially on digital media. Digital advertising and social media engagement are increasingly being used to promote UPFs. These platforms enable targeted advertising that can significantly influence consumer behavior (Al-Ababneh et al., 2025). In addition, many people, especially children and adolescents, tend to snack while watching a screen. Snacks consumed while using screens are frequently UPF, which are convenient and tasty, but unhealthy (Rodríguez-Barniol et al., 2024).

Parental attitudes and control over food choices also play an essential role in shaping this behavior (Rodríguez-Barniol et al., 2024). Parental behavior and the home environment play a significant role in shaping adolescents' eating habits. In this study, participants who lived with only one parent (father or mother) or did not live with both parents had higher energy intake from UPFs. Adolescents living with both parents are less likely to consume UPFs compared to those living with a single parent or without any parents (Souza et al., 2023). Parents play a significant

Table 3. Determining factors of the domination of energy intake in adolescent girls in Tasikmalaya City

Determinant of UPF	Domination of energy intake				<i>p</i> ; cOR (95% CI)
	UPF		Non-UPF		
	n	%	n	%	
Individual factors					
Age (years)					
12-13	52	22.9	175	77.1	0.593; 1.148 (0.753-1.751)
14-15	59	20.6	228	79.4	1.00
Grade					
VII	34	19.0	145	81.0	0.350; 0.786 (0.500-1.235)
VIII-IX	77	23.0	258	77.0	1.00
Pocket money (IDR per day)					
> 20,000	22	22.9	74	77.1	0.833; 1.099 (0.647-1.868)
≤ 20,000	89	21.3	329	78.7	1.00
Sleep duration (hours per day)					
< 8	63	20.6	243	79.4	0.573; 0.864 (0.656-1.322)
≥ 8	48	23.1	160	76.9	1.00
Late-night sleeping					
Yes	17	27.4	45	72.6	0.306; 1.439 (0.788-2.628)
No	95	20.8	358	79.2	1.00
Screen time duration (hours per day)					
> 4	91	25.7	263	74.3	0.001*; 2.422 (1.432-4.097)
≤ 4	20	12.5	140	87.5	1.00
Physical activity (times per week)					
< 3	98	22.5	338	77.5	0.318; 1.450 (0.767-2.740)
≥ 3	13	16.7	65	83.3	1.00
Nutrition literacy					
Bad	64	24.4	198	75.6	0.138; 1.410 (0.923-2.155)
Good	47	18.7	205	81.3	1.00
Main meals (times per day)					
< 3	94	24.8	285	75.2	0.005*; 2.289 (1.309-4.005)
≥ 3	118	12.6	135	87.4	1.00
Breakfast habits					
Rare	72	26.9	196	73.1	0.003*; 1.950 (1.261-3.015)
Frequent	39	15.9	207	84.1	1.00
Interpersonal factors					
Father's education					
Elementary-Junior high school	37	19.6	152	80.4	0.461; 0.745 (0.530-1.286)
Senior high school-university	74	22.8	251	77.2	1.00
Mother's education					
Elementary-Junior high school	50	25.0	150	75.0	0.165; 1.383 (0.904-2.115)
Senior high school-university	61	19.4	253	80.6	1.00
Father's occupation					
Not working	4	20.0	16	80.0	1.000; 0.904 (0.296-2.761)
Working	107	21.7	387	78.3	1.00
Mother's occupation					
Working	37	27.6	97	72.4	0.065; 1.577 (1.000-2.488)
Not working	72	19.5	306	80.5	1.00
Family members (persons)					
> 4	54	23.0	181	77.0	0.554; 1.164 (0.763-1.770)
≤ 4	57	20.4	222	79.6	1.00

Table 3. *Continued*

Determinant of UPF	Domination of energy intake				<i>p</i> ; cOR (95% CI)
	UPF		Non-UPF		
	n	%	n	%	
Family status					
Incomplete	19	33.3	38	66.7	0.035*; 1.984 (1.093-3.601)
Complete	92	20.1	365	79.9	1.00

Note: *significant at $p < 0.05$; cOR = Crude odd ratio; 1.00 as reference

Table 4. Final model logistic regression of determinants of energy intake from UPFs in adolescent girls in Tasikmalaya City

Variable	<i>p</i> -Value	aOR	95% CI
Screen time	0.001	2.424	1.420-4.140
Breakfast habits	0.094	1.466	0.936-2.295
Main meals	0.019	1.996	1.121-3.554
Family status	0.006	2.336	1.275-4.282

Note: aOR = Adjusted odds ratio

role in shaping adolescents' eating patterns through various mechanisms, including serving as role models (Pitt and Curran, 2023), influencing the home food environment, and establishing mealtime rules (Souza et al., 2023).

In this study, participants who had fewer than 3 main meals per day had higher energy intake from UPFs than from non-UPFs. There has been a decline in the habit of having 3 main meals a day, particularly among women, with a concurrent increase in UPFs consumption (Ribeiro et al., 2023). Studies show that reducing meal frequency (e.g., consuming fewer than 3 meals per day) can lead to a significant increase in appetite and decreased satiety. This suggests that fewer meals can make a person feel hungrier and more susceptible to food cravings. This aligns with findings that reducing meal frequency can increase food cravings due to the body's response to perceived nutrient deficiencies (Meule, 2020).

Food cravings are closely related to high-calorie food consumption. Individuals with high food cravings (HC) are more likely to think about and consume high-calorie snacks compared to those with low food cravings (LC) (Devoto et al., 2025). Furthermore, high-calorie food cravings predict their consumption, which contributes to unhealthy eating habits and potential weight gain (Werthmann et al., 2023). In general, people who eat less than 3 main meals per day will miss one of the standard meal times, namely breakfast, lunch, or dinner (Sun et al., 2023). Adolescents who skipped one of the main meals, such as breakfast, experienced an increase in average

energy intake from UPFs, along with increases in BMI-for-age z-scores (BAZ) and body fat percentage (Cândido et al., 2024). Skipping breakfast has been shown to increase hunger and appetite throughout the day, which can lead to overeating at the next meal, such as lunch (Fatima et al., 2020). Skipping meals is a driving factor in unhealthy eating habits, including increased consumption of UPFs among adolescents. This is because UPFs are often more accessible and appealing to adolescents seeking quick, easy meal options (Saleh and Maala, 2021). In this study, participants who ate fewer than 3 main meals per day had higher energy consumption from UPFs than from non-UPFs.

Participants of this study were junior high school students, with study time at school for 8 hours (starting from 7.00 AM to 2.00 PM) (Ministry of Education and Culture of Indonesia, 2017). Therefore, students are still at school during lunchtime. Students who eat breakfast will feel hungry around 12 noon. This is because the average gastric emptying time after eating is about 5.8 ± 0.8 hours, with a significant decrease in gastric contents observed around 4 hours after eating (Cho et al., 2019). To satisfy their hunger, they will buy food provided by the school. In Indonesia, school canteens often provide unhealthy food and drinks, including fried snacks and sweet beverages (Rachmadewi et al., 2021), with low nutritional quality (Sijangga et al., 2026). In addition, various types of food and drinks in the UPFs category are widely sold in school canteens in Indonesia (Ginting et al.,

2024). UPF is often offered in school canteens, including sausages, packaged snacks, sweet drinks, and candy. For example, in Brazilian schools, the presence of canteens is associated with a higher frequency of consumption of these foods among adolescents (Leite et al., 2021). School canteens often sell UPFs, which are industrially formulated products that are typically high in sugars, fats, and salts, and low in essential nutrients. These foods are designed to be convenient, tasty, and appealing, making them popular among students (Leffa, 2023).

CONCLUSIONS

This study revealed that in more than one-fifth of the participants, energy intake from UPFs was more dominant than that from non-UPFs. A socioecological approach was used to describe the determinants of energy intake from UPFs. Individual factors, including screen time of > 4 hours per day and consuming main meals < 3 times a day, are risk factors for increased energy intake from UPFs. Meanwhile, living with an incomplete family is an interpersonal factor that increases energy intake from UPFs. Parents must monitor their children's nutritional intake, especially by limiting UPF consumption. Policies regarding healthy snacks sold in school canteens also need to be implemented.

ACKNOWLEDGEMENTS

The authors thank all the students who participated in this research. The authors also thank the LPPM Universitas Siliwangi for funding this research through Rector's Decree No. 1111/UN58/P/2025.

REFERENCES

- Al-Ababneh, H. A., Alzyadat, A. Y., Al-Husban, A. M., Alotoum, F. J., Al Kurdi, B., Ibragimkhalilova, T., & Barhoush, F. M. S. (2025). Marketing in social networks: Key trends and development forecasts. *International Review of Management and Marketing*, 15(4), 388–396. <https://doi.org/10.32479/irmm.19195>
- Andarwulan, N., Madanijah, S., Briawan, D., Anwar, K., Bararah, A., Saraswati, & Średnicka-Tober, D. (2021). Food consumption pattern and the intake of sugar, salt, and fat in the South Jakarta City—Indonesia. *Nutrients*, 13(4), 1289. <https://doi.org/10.3390/nu13041289>
- Andrade, G. C., Gombi-Vaca, M. F., Louzada, M. L. D. C., Azeredo, C. M., & Levy, R. B. (2020). The consumption of ultra-processed foods according to eating out occasions. *Public Health Nutrition*, 23(6), 1041–1048. <https://doi.org/10.1017/S1368980019002623>
- Bahramfard, T., Omid Salehi, S., Akbartabar Toori, M., Pourmahmoudi, A., Jowshan, M., Parvin, S., ... & Malekzadeh, J. (2020). Nutritional literacy status and its related factors in students of Yasuj University of Medical Sciences. *Nutrición Clínica y Dietética Hospitalaria*, 40(4), 55–62. <https://doi.org/10.12873/404malekzadeh>
- Bestari, F., Andarwulan, N., & Palupi, E. (2023). Synthesis of effect sizes on dose response from ultra-processed food consumption against various non-communicable diseases. *Foods*, 12(24), 4457. <https://doi.org/10.3390/foods12244457>
- Cândido, A. C. de O., Neves, F. S., Fontes, V. S., Melo, A. S. T., de Faria, E. R., Netto, M. P., ... & Cândido, A. P. C. (2024). Frequency of breakfast consumption and its associations with food consumption by degree of industrial processing and with indicators of overweight in Brazilian adolescents (EVA-JF Study). *Nutrition*, 117, 112226. <https://doi.org/10.1016/j.nut.2023.112226>
- CDC. (2020). *National health and nutrition examination survey: Questionnaires, datasets, and related documentation*. Centers for Disease Control and Prevention, *National Center for Health Statistics*. Retrieved from <https://www.cdc.gov/nchs/nhanes/default.aspx>
- Cho, E. A., Kim, M. S., Cha, Y. B., Lee, M. S., & Song, T. (2019). Evaluation of gastric emptying time of a rice-based meal using serial sonography. *BioMed Research International*, 2019(1), 5917085. <https://doi.org/10.1155/2019/5917085>
- Choudhury, S., Bi, A. Z., Medina-Lara, A., Morrish, N., & Veettil, P. C. (2025). The rural food environment and its association with diet, nutrition status, and health outcomes in low-income and middle-income countries

- (LMICs): A systematic review. *BMC Public Health*, 25(1), 994. <https://doi.org/10.1186/s12889-025-22098-4>
- Colozza, D., Wang, Y. C., & Avendano, M. (2023). Does urbanisation lead to unhealthy diets? Longitudinal evidence from Indonesia. *Health & Place*, 83, 103091. <https://doi.org/10.1016/j.healthplace.2023.103091>
- Cortes, C., Brandão, J. M., Cunha, D. B., Paravidino, V. B., & Sichieri, R. (2023). Blood pressure variation and ultra-processed food consumption in children with obesity. *European Journal of Pediatrics*, 182(9), 4077–4085. <https://doi.org/10.1007/s00431-023-05076-z>
- Costa, R. M., Oliveira, A. G., Torres, K. G., Souza, A. M., Pereira, G. S., & Bezerra, I. W. L. (2024). Quantitative assessment of the inadequate intake of macronutrients, minerals, and vitamins associated with ultra-processed food consumption. *International Journal of Environmental Research and Public Health*, 21(7), 888. <https://doi.org/10.3390/ijerph21070888>
- Devoto, F., Girati, M., Asteria, C., Ravelli, M., Paulesu, E., & Zapparoli, L. (2025). Satiety and food availability modulate the approach and inhibitory responses to food cues in healthy weight participants. *Appetite*, 214, 108166. <https://doi.org/10.1016/j.appet.2025.108166>
- Dewanti, R., Probandari, A., & Mulyani, S. (2022). Factors affecting the choices of healthy diet among adolescents in rural areas. *Media Gizi Indonesia*, 17(1), 204–212. <https://doi.org/10.204736/mgi.v17i1.204-212>
- Fatima, N., Noor, Z., & Shafique, S. (2020). Effect of breakfast skipping on body mass index in adolescent university girls. *Journal of University Medical and Dental College*, 11(2), 9–15. <https://doi.org/10.37723/jumdc.v11i2.407>
- Ginting, S. M. D., Februhartanty, J., & Khusun, H. (2024). Association between consumption of ultra-processed foods and beverages with nutritional status of private senior high school students in Pontianak, West Kalimantan, Indonesia. *World Nutrition Journal*, 7(i2), 90–103. <https://doi.org/10.25220/wnj.v07.i2.0011>
- Gonçalves, H. V. B., Batista, L. S., de Amorim, A. L. B., & Bandoni, D. H. (2023). Association between consumption of ultra-processed foods and sociodemographic characteristics in Brazilian adolescents. *Nutrients*, 15(9), 2027. <https://doi.org/10.3390/nu15092027>
- Grasaas, E., Ostojic, S., & Jahre, H. (2024). Adherence to sleep recommendations is associated with higher satisfaction with life among Norwegian adolescents. *BMC Public Health*, 24(1), 1288. <https://doi.org/10.1186/s12889-024-18725-1>
- Gutkowska, K., Wierzbicka, E., Madej, D., Czarniecka-Skubina, E., & Hamulka, J. (2025). Breakfast frequency, lifestyle-related factors and their association with body weight status among Polish primary school children aged 10 to 12 years: Results from a nationwide cross-sectional study. *Nutrition Journal*, 24(1), 160. <https://doi.org/10.1186/s12937-025-01231-4>
- Hidayanti, L., Saraswati, D., & Aisyah, I. S. (2023). Food intake, dietary quality and nutritional status of female adolescent in Tasikmalaya, West Java, Indonesia. *Malaysian Journal of Medicine and Health Sciences*, 19(6), 95–100. <https://doi.org/10.47836/mjmhs.19.6.13>
- Huang, K., Zhao, L., Guo, Q., Yu, D., Yang, Y., Cao, Q., ... & Fang, H. (2022). Comparison of the 24 h dietary recall of two consecutive days, two non-consecutive days, three consecutive days, and three non-consecutive days for estimating dietary intake of Chinese adult. *Nutrients*, 14(9), 1960. <https://doi.org/10.3390/nu14091960>
- Ishikawa-Takata, K., Kurotani, K., Adachi, M., & Takimoto, H. (2021). Frequency of meals that includes staple, main and side dishes and nutrient intake: Findings from the 2012 National Health and Nutrition Survey, Japan. *Public Health Nutrition*, 24(9), 2618–2628. <https://doi.org/10.1017/S1368980020001202>
- Islam, M. R., Rahman, S. M., Rahman, M. M., Pervin, J., Rahman, A., & Ekström, E. C. (2022). Gender and socio-economic stratification of ultra-processed and deep-fried food consumption among rural adolescents: A cross-sectional study from Bangladesh.

- PLoS ONE*, 17(7), e0272275. <https://doi.org/10.1371/journal.pone.0272275>
- James, J., Pringle, A., Mourton, S., & Roscoe, C. M. P. (2023). The effects of physical activity on academic performance in school-aged children: A systematic review. *Children*, 10(6), 1019. <https://doi.org/10.3390/children10061019>
- Kesaite, V., Chavez-Ugalde, Y., White, M., & Adams, J. (2024). Overlap between ultra-processed food and food that is high in fat, salt or sugar: Analysis of 11 annual waves of the UK National Diet and Nutrition Survey 2008/09-2018/19. *medRxiv*, 2024-08. <https://doi.org/10.1101/2024.08.27.24312650>
- Laksono, A. D., & Wulandari, R. D. (2021). The factors correlate to family size in Indonesia. *Aspirasi: Jurnal Masalah-Masalah Sosial*, 12(1), 1–13. <https://doi.org/10.46807/aspirasi.v12i1.2066>
- Lane, M. M., Gamage, E., Du, S., Ashtree, D. N., McGuinness, A. J., Gauci, S., ... & Marx, W. (2024). Ultra-processed food exposure and adverse health outcomes: Umbrella review of epidemiological meta-analyses. *BMJ*, 384, e077310. <https://doi.org/10.1136/bmj-2023-077310>
- Leffa, P. (2023). Ultra-processed foods. *Encyclopedia of Human Nutrition (Fourth Edition)* (pp. 653–662). <https://doi.org/10.1016/B978-0-12-821848-8.00009-3>
- Leite, M. A., Azeredo, C. M., Peres, M. F. T., Escuder, M. M. L., & Levy, R. B. (2021). Availability and consumption of ultra-processed foods in schools in the municipality of Sao Paulo, Brazil: Results of the SP-Proso. *Cadernos de Saude Publica*, 37, e00162920. <https://doi.org/10.1590/0102-311X00162920>
- Li, H., Wang, Y., Sonestedt, E., & Borné, Y. (2023). Associations of ultra-processed food consumption, circulating protein biomarkers, and risk of cardiovascular disease. *BMC Medicine*, 21(1), 415. <https://doi.org/10.1186/s12916-023-03111-2>
- Martini, D., Godos, J., Bonaccio, M., Vitaglione, P., & Grosso, G. (2021). Ultra-processed foods and nutritional dietary profile: A meta-analysis of nationally representative samples. *Nutrients*, 13(10), 3390. <https://doi.org/10.3390/nu13103390>
- Matos, R. A., Adams, M., & Sabaté, J. (2021). Review: The consumption of ultra-processed foods and non-communicable diseases in Latin America. *Frontiers in Nutrition*, 8, 622714. <https://doi.org/10.3389/fnut.2021.622714>
- McLeroy, K. R., Bibeau, D., Steckler, A., & Glanz, K. (1988). An ecological perspective on health promotion programs. *Health Education Quarterly*, 15(4), 351–377. <https://doi.org/10.1177/109019818801500401>
- Mensah, J. P., Akparibo, R., Thomas, C., Aryeetey, R., & Brennan, A. (2025). Application of the socioecological model to understand the drivers of excessive alcohol and salt consumption: A qualitative study in Ghana. *BMJ Open*, 15(10), e100490. <https://doi.org/10.1136/bmjopen-2025-100490>
- Mescoloto, S. B., Pongiluppi, G., & Domene, S. M. Á. (2024). Ultra-processed food consumption and children and adolescents' health. *Jornal de Pediatria*, 100(Supplement 1), S18–S30. <https://doi.org/10.1016/j.jpmed.2023.09.006>
- Meule, A. (2020). The psychology of food cravings: The role of food deprivation. *Current Nutrition Reports*, 9(3), 251–257. <https://doi.org/10.1007/s13668-020-00326-0>
- Ministry of Education and Culture of Indonesia. (2017). *Regulation of the Minister of Education and Culture Number 23 of 2017 concerning School Days*. Retrieved from <https://peraturan.go.id/id/permendikbud-no-23-tahun-2017>
- Ministry of Health of Indonesia. (2023). *Annual Report 2022: Prevention and Control of Non-Communicable Diseases (NCDs)*.
- Monteiro, C. A., Cannon, G., Lawrence, M., Costa Louzada, M. D., & Pereira Machado, P. (2019). *Ultra-processed foods, diet quality, and health using the NOVA classification system*. Rome: FAO, 48. Retrieved from https://openknowledge.fao.org/server/api/core/bitstreams/5277b379-0acb-4d97-a6a3-602774104629/content?utm_source=coast%20reporter&utm_campaign=coast%20reporter%3A%20outbound&utm_medium=referral

- Monteiro, C. A., Louzada, M. L., Steele-Martinez, E., Cannon, G., Andrade, G. C., Baker, P., ... & Touvier, M. (2025). Ultra-processed foods and human health: The main thesis and the evidence. *The Lancet*, 406(10520), 2667–2684. [https://doi.org/10.1016/S0140-6736\(25\)01565-X](https://doi.org/10.1016/S0140-6736(25)01565-X)
- Moodie, R., Bennett, E., Kwong, E. J. L., Santos, T. M., Pratiwi, L., Williams, J., & Baker, P. (2021). Ultra-processed profits: The political economy of countering the global spread of ultra-processed foods - A synthesis review on the market and political practices of transnational food corporations and strategic public health responses. *International Journal of Health Policy and Management*, 10(12), 968–982. <https://doi.org/10.34172/ijhpm.2021.45>
- Moradi, S., Kermani, M. A. H., Bagheri, R., Mohammadi, H., Jayedi, A., Lane, M. M., ... & Suzuki, K. (2021). Ultra-processed food consumption and adult diabetes risk: A systematic review and dose-response meta-analysis. *Nutrients*, 13(12), 4410. <https://doi.org/10.3390/nu13124410>
- Mukti, B. H. (2025). Sample size determination: Principles and applications for health research. *Health Sciences International Journal*, 3(1), 127–143. <https://doi.org/10.71357/hsij.v3i1.63>
- Muraro, A. P., Lobato, J. C. P., Rodrigues, P. R. M., & Souza, A. M. (2023). Dietary sources of energy, saturated fat, trans fat, and added sugars in Brazilian adolescents. *Nutrire*, 48(1), 27. <https://doi.org/10.1186/s41110-023-00213-y>
- Nurhasan, M., Ariesta, D. L., Utami, M. M. H., Fahim, M., Aprillyana, N., Maulana, A. M., & Ickowitz, A. (2024). Dietary transitions in Indonesia: The case of urban, rural, and forested areas. *Food Security*, 16(6), 1313–1331. <https://doi.org/10.1007/s12571-024-01488-3>
- Ouvrein, G., Verswijvel, K., & Kimpe, D. L. (2019). Social goals in context: Social media. *Social Goals in the Classroom: Findings on Student Motivation and Peer Relations* (pp. 237–254). <https://doi.org/10.4324/9780429468452-14>
- Pereira, R. B., Martini, T. C., Buchweitz, C., Kieling, R. R., Fisher, H. L., Kohrt, B. A., ... & Kieling, C. (2024). Self-reported social media use by adolescents in Brazil: A school-based survey. *Trends in Psychiatry and Psychotherapy*, 46, e20220545. <https://doi.org/10.47626/2237-6089-2022-0545>
- Pitt, P. D., & Curran, K. A. (2023). Like mother, like son: An adolescent who diets with his parent. *Fad Diets and Adolescents: A Guide for Clinicians, Educators, Coaches and Trainers* (pp. 83–87). https://doi.org/10.1007/978-3-031-10565-4_11
- Pradana, M. A., Farapti, F., & Daud, Z. A. M. (2025). Analysis of sodium content in commercially salty snack and its contribution to daily salt intake: A market survey in East Surabaya. *Amerta Nutrition Journal*, 9(3), 538–544. Retrieved from <https://e-journal.unair.ac.id/AMNT/upcoming/view/70102>
- Provincial Government of West Java. (2024). *Strengthening Commitment on Decade of Actions Toward Resilient West Java to Leave No One Behind*. Retrieved from <https://www.iges.or.jp/en/vlr/west-java>
- Putri, A. A., & Anwar, K. (2024). The relationship of energy and nutrient intake from eating out and eating at home on blood hemoglobin levels in adolescents. *Indonesian Journal of Public Health Nutrition*, 4(2), 8209. <https://doi.org/10.7454/ijphn.v4i2.8209>
- Rachmadewi, A., Soekarjo, D., Maehara, M., Alwi, B., Mulati, E., & Rah, J. H. (2021). School canteens in selected areas in Indonesia: A situation analysis. *Food and Nutrition Bulletin*, 42(2), 225–246. <https://doi.org/10.1177/03795721211008021>
- Raihandhany, R., & Purnomo. (2025). Quantitative ethnobotany approach on local food plants in Banceuy Traditional Village, Subang Regency, West Java. *Media Konservasi*, 30(1), 172–183. <https://doi.org/10.29244/medkon.30.1.172>
- Republic of Indonesia. (2003). *Law of the Republic of Indonesia Number 20 of 2003 Concerning The National Education System*. Jakarta. Retrieved from <https://pusmendik.kemdikbud.go.id/pdf/file-154>

- Ribeiro, G. J., Mendes, A. E. P., Costa, E. D. A., & Carvalho, D. V. (2023). Trends in food consumption by adults in a Brazilian northeastern state. *Journal of Taibah University Medical Sciences*, 18(6), 1261–1267. <https://doi.org/10.1016/j.jtumed.2023.04.008>
- Rocha, L. L., Gratão, L. H. A., Carmo, A. S. D., Costa, A. B. P., Cunha, C. D. F., Oliveira, T. R. P. R. D., & Mendes, L. L. (2021). School type, eating habits, and screen time are associated with ultra-processed food consumption among Brazilian adolescents. *Journal of the Academy of Nutrition and Dietetics*, 121(6), 1136–1142. <https://doi.org/10.1016/j.jand.2020.12.010>
- Rodríguez-Barniol, M., Pujol-Busquets, G., & Bach-Faig, A. (2024). Screen time use and ultra-processed food consumption in adolescents: A focus group qualitative study. *Journal of the Academy of Nutrition and Dietetics*, 124(10), 1336–1346. <https://doi.org/10.1016/j.jand.2024.04.015>
- Saleh, B. B., & Maala, E. G. (2021). Impact of adolescents' life style and reasons of skipping essential meals upon their BMI at secondary schools in Baghdad City. *Indian Journal of Forensic Medicine and Toxicology*, 15(1), 2507–2514. <https://doi.org/10.37506/ijfmt.v15i1.13777>
- Sandoval-insausti, H., Jim, M., Donat-vargas, C., Rey-garc, J., Banegas, J. R., Rodr, F., & Castillon, P. G. (2020). Ultra-processed food consumption is associated with abdominal obesity: A prospective cohort study in older adults. *Nutrients*, 12(8), 2368. <https://doi.org/10.3390/nu12082368>
- Santos, M., & Assunção, R. (2025). Food choice, nutrition, and public health. *Foods*, 14(7), 1243. <https://doi.org/10.3390/foods14071243>
- Shinde, S., Harling, G., Assefa, N., Bärnighausen, T., Bukenya, J., Chukwu, A., ... & Fawzi, W. (2023). Counting adolescents in: The development of an adolescent health indicator framework for population-based settings. *EClinicalMedicine*, 61, 102067. <https://doi.org/10.1016/j.eclinm.2023.102067>
- Shim, J. S., Shim, S. Y., Cha, H. J., Kim, J., & Kim, H. C. (2022). Association between ultra-processed food consumption and dietary intake and diet quality in Korean adults. *Journal of the Academy of Nutrition and Dietetics*, 122(3), 583–594. <https://doi.org/10.1016/j.jand.2021.07.012>
- Sijangga, M. O., Hositanisita, H., Lewis, E. C., Hadi, H., Matsuzaki, M., Surkan, P. J., ... & Gittelsohn, J. (2026). Cross-sectional study of prepared foods sold in Indonesian school canteens to inform childhood obesity programs and policies. *Journal of Nutritional Science*, 15, e7. <https://doi.org/10.1017/jns.2025.10068>
- Singh, J. A., Siddiqi, M., Parameshwar, P., & Chandra-Mouli, V. (2019). World Health Organization Guidance on ethical considerations in planning and reviewing research studies on sexual and reproductive health in adolescents. *Journal of Adolescent Health*, 64(4), 427–429. <https://doi.org/10.1016/j.jadohealth.2019.01.008>
- Sparrow, R., Agustina, R., Bras, H., Sheila, G., Rieger, M., Yumna, A., ... & Melse-Boonstra, A. (2021). Adolescent nutrition—Developing a research agenda for the second window of opportunity in Indonesia. *Food and Nutrition Bulletin*, 42(1_suppl), S9–S20. <https://doi.org/10.1177/0379572120983668>
- Sun, Y., Rong, S., Liu, B., Du, Y., Wu, Y., Chen, L., & Bao, W. (2023). Meal skipping and shorter meal intervals are associated with increased risk of all-cause and cardiovascular disease mortality among US adults. *Journal of the Academy of Nutrition and Dietetics*, 123(3), 417–426.e3. <https://doi.org/10.1016/j.jand.2022.08.119>
- Souza, T. N., Andrade, G. C., Rauber, F., Levy, R. B., & Da Costa Louzada, M. L. (2022). Consumption of ultra-processed foods and the eating location: Can they be associated? *British Journal of Nutrition*, 128(8), 1587–1594. <https://doi.org/10.1017/S0007114521004992>
- Souza, M. R. D., De Souza Andrade, A. C., Froelich, M., Muraro, A. P., & Rodrigues, P. R. M. (2023). Association of household composition with dietary patterns among adolescents in Brazil. *British Journal of Nutrition*, 130(7), 1213–1219. <https://doi.org/10.1017/S000711452300020X>

- Thapsuwan, S., Phulkerd, S., Chamrathirong, A., Gray, R. S., Jindaratnaporn, N., Loyfah, N., ... & Pattaravanich, U. (2024). Relationship between consumption of high fat, sugar or sodium (HFSS) food and obesity and non-communicable diseases. *BMJ Nutrition, Prevention and Health*, 7(1), 78–87. <https://doi.org/10.1136/bmjnph-2023-000794>
- UNICEF Indonesia. (2021). *Social and behaviour change communication strategy: Improving adolescent nutrition in Indonesia*. United States of America. Retrieved from <https://coilink.org/20.500.12592/49thqg0>
- United Nations. (2019). *World population prospects 2019 highlights*. New York. Retrieved from https://population.un.org/wpp/assets/Files/WPP2019_Highlights.pdf
- Vashtianada, A., Setiarini, A., & Sartika, R. A. D. (2023). The difference of ultra-processed food consumption based on individual characteristics and other factors among non-health undergraduate students in Universitas Indonesia in 2023. *Indonesian Journal of Public Health Nutrition*, 4(1), 59–71. <https://doi.org/10.7454/ijphn.v4i1.7393>
- Vitale, M., Costabile, G., Testa, R., D'Abbronzio, G., Nettore, I. C., Macchia, P. E., & Giacco, R. (2024). Ultra-processed foods and human health: A systematic review and meta-analysis of prospective cohort studies. *Advances in Nutrition*, 15(1), 100121. <https://doi.org/10.1016/j.advnut.2023.09.009>
- Wellard-Cole, L., Davies, A., Chen, J., Jung, J., Bente, K. B., Kay, J., ... & Allman-Farinelli, M. (2021). The contribution of foods prepared outside the home to the diets of 18- to 30-year-old australians: The mymeals study. *Nutrients*, 13(6), 1761. <https://doi.org/10.3390/nu13061761>
- Werthmann, J., Tuschen-Caffier, B., Ströbele, L., Kübel, S. L., & Renner, F. (2023). Healthy cravings? Impact of imagined healthy food consumption on craving for healthy foods and motivation to eat healthily-Results of an initial experimental study. *Appetite*, 183, 106458. <https://doi.org/10.1016/j.appet.2023.106458>
- Wang, M., Du, X., Huang, W., & Xu, Y. (2022). Ultra-processed foods consumption increases the risk of hypertension in adults: A systematic review and meta-analysis. *American Journal of Hypertension*, 35(10), 892–901. <https://doi.org/10.1093/ajh/hpac069>
- Yetmi, F., Harahap, F. S. D., & Lestari, W. (2021). Analisis faktor yang memengaruhi konsumsi fast food pada siswa di SMA Cerdas Bangsa Kabupaten Deli Serdang tahun 2020. *Studia: Jurnal Hasil Penelitian Mahasiswa*, 6(1), 24–47. <https://doi.org/10.32923/stu.v6i1.2021>
- Zablotsky, B., Ng, A. E., Black, L. I., Haile, G., Bose, J., Jones, J. R., & Blumberg, S. J. (2025). Associations between screen time use and health outcomes among US teenagers. *Preventing Chronic Disease*, 22, E38. <https://doi.org/10.5888/pcd22.240537>
- Zahra, S. A. (2023). *Strategi integrated marketing communication rumah potong ayam (rpa) CV Sukahati Pratama Kota Tasikmalaya*. Universitas Islam Indonesia. Retrieved from <https://dspace.uui.ac.id/123456789/55651>
- Zhong, C. C., Chen, M., Li, Z., Yang, Z., Keung, V. M. W., Lo, A., ... & Wong, M. C. S. (2025). Examining sleep habits and associated lifestyle factors in adolescents: A population-based study. *Brain and Behavior*, 15(10), e70885. <https://doi.org/10.1002/brb3.70885>