

# Agrotechnology Research Journal Volume 9, No. 1, June 2025, pp. 52–xx

ISSN 2655-7924 (Print) ISSN 2614-7416 (Online) https://jurnal.uns.ac.id/arj doi:10.20961/agrotechresj.v9i1.102718

# Potential Decline in Domestic Rice Production Due to The Degradation of Land Functions from Paddy Fields to Other Uses: A Case Study in Banyumas Regency, Central Java, Indonesia

Bondansari Bondansari¹⁺, Widiatmaka Widiatmaka², Machfud Machfud³, Khursatul Munibah², Wiwin Ambarwulan⁴

<sup>1</sup>Department of Agrotechnology, Faculty of Science and Technology, Universitas Nahdlatul Ulama, Banyumas, Central Java 53145, Indonesia

Received May 30, 2025; Accepted June 15, 2025

#### **ABSTRACT**

Shallot cv. Tajuk is one of the most preferred shallot varieties to be cultivated, especially by farmers living in lowland areas with low water availability issues, such as Bantul and Kulon Progo. However, to date, the information about the resistance of this variety to Fusarium disease is poorly studied. Therefore, this study aims to investigate the prevalence of Fusarium disease in shallot cv. Tajuk is cultivated in Bantul and Kulon Progo. A survey of Fusarium disease occurrence was conducted in 8 shallot fields from 3 districts in Bantul (Imogiri, Sanden, and Kretek) and six shallot fields representing two districts of Kulon Progo (Lendah and Sentolo). Disease observation in Bantul was carried out during the dry season from June to August 2023, while the survey in Kulon Progo was performed during the rainy season from January to February 2024. The cropping system applied in each selected shallot field was also collected through the landowner interviews. Data obtained from disease monitoring were statistically analyzed using one-way analysis of variance (ANOVA), and significance among locations was subsequently analyzed using Duncan's New Multiple Range Test with a *p*<0.05. Findings of this study revealed that the susceptibility of shallot cv. Tajuk to Fusarium disease varied between Bantul and Kulon Progo. Shallot var. Tajuk conferred higher resistance against Fusarium disease when it was cultivated in Kulon Progo. Results of this study also confirmed that different farming practices applied in each location appear to have a minor impact on the severity of this disease in the shallot cv. Tajuk.

Keywords: Allium cepa; Disease occurrence; Leaf twisting disease; Resistance; Variety

*Cite this as (CSE Style)*: Aisyah SN, Fajrin QV, Dewi IP, Hidayat T, Astuti A, Harikrishna JA. 2025. Susceptibility of *Shallot* cv Tajuk cultivated in Bantul and Kulon Progo against *Fusarium* disease. Agrotechnology Res J. 9(1):44–51. https://dx.doi.org/10.20961/agrotechresj.v9i1.105240.

### INTRODUCTION

Shallot is considered one of the strategic horticultural crops that contributes significantly to national economic growth. High demand and consumption of shallots in Indonesia are associated with its use as a basic ingredient for cooking (Anggrasari et al. 2025; Ridqi and Kusumaningsih 2025). Apart from its use as food flavoring, latest studies also reported the promising medicinal properties of shallot (Gusman et al. 2025; Neda et al. 2025). Moreover, Ridqi and Kusumaningsih (2025) mentioned that shallot has high export potential and high employment opportunities. Due to its economic value, shallots have become a commodity that could

affect the inflation rate in Indonesia, suggesting their essential contribution to national food security (Anggrasari et al. 2025; Arifin et al. 2025).

The special Region of Yogyakarta is one of the shallot-producing provinces in Indonesia (Anggrasari et al. 2025) that contributed about 333,117 quintals the shallot production in 2024 (Statistics Indonesia 2025). Shallot production in this province is centralized in two regencies, namely Bantul and Kulon Progo. Considering the size of the harvested area, Bantul Regency is the largest shallot-producing area with the harvested area in 2024, reaching 1,806 ha and total production of about 198,286 quintals. Shallot production in Kulon Progo Regency was 128,345 quintals in 2024 with the harvested area reaching 983.2 ha (Statistics Indonesia 2025). According to these data, the shallot productivity in these two regions is approximately 10.99 t.ha-1 (Bantul) and 13.04 t.ha-1 (Kulon Progo). This productivity is

\*Corresponding Author: E-Mail: sitinur@umy.ac.id



<sup>&</sup>lt;sup>2</sup>Department of Soil Science and Land Resources, Institut Pertanian Bogor, Bogor, West Java 16680, Indonesia.

<sup>&</sup>lt;sup>3</sup>Department of Agricultural Industrial Technology, Institut Pertanian Bogor, West Java 16680 , Indonesia

<sup>&</sup>lt;sup>5</sup>National Research and Innovation Agency, Central Jakarta, Jakarta 10340, Indonesia

considered low compared to the world average productivity (21.4 t.ha-1) (FAOSTAT 2023). This low productivity suggests that the shallot cultivation in these regions was challenging. Sutardi et al. (2022) highlighted low efficiency farming practices as one of main factors limiting the shallot productivity, particularly in Special Region of Yogyakarta. Moreover, Arifin et al. (2025) also explained the challenges faced by shallot farmers in Bantul Regency due to unstable weather and disease infection.

Regarding disease infection, soil-borne disease caused by Fusarium spp. has been reported as an important disease in the production of Allium species, including shallot (Poursakhi et al. 2024). The disease is locally as *moler* disease and known internationally as basal plate rot or basal rot. When shallots are infected by this disease, various symptoms may appear, such as leaf twisting followed by withering, root rot, bulb rot, and basal plate rot (Le et al. 2021; Sholeh and Nurcahyanti 2023). The importance of Fusarium disease in the cultivation of Allium species has been extensively documented due to its challenging management, thus leading to significant crop loss (Dewi et al. 2025; Haapalainen et al. 2023; Sharma et al. 2024). Infection of Fusarium disease had caused yield loss in shallots up to 78% and severe damage to stored bulbs (Dinata et al. 2021; Mariani et al. 2022). Previous studies also reported significant crop loss up to 50% in onion caused by Fusarium infection (Mahanty et al. 2023; Sharma et al. 2024; Sogoba et al. 2022).

As one of the shallot-producing provinces in Indonesia, the Special Region of Yogyakarta cultivates several varieties of shallots, including Tajuk variety. This variety, also known as the Thailand variety, has been widely cultivated due to its suitability for lowlands and its high adaptability with minimal water input (Fauzia et al. 2020; Pasaribu 2017). Moreover, Pasaribu (2017) also highlighted other advantages of shallot cv. Tajuk, such as the ability to grow in acidic soil (pH 5.5), tiller

production ranging from 5-15 per plant, and productivity of about 10 t.ha-1 dried bulb. Despite its promising potential, the information regarding its susceptibility to Fusarium disease is not yet thoroughly studied to date. Concerning farmers' preferences regarding this variety, the investigation of Fusarium disease occurrence in shallot cv. Tajuk has been urgently needed. Therefore, this present study aimed to investigate the prevalence of Fusarium disease during the cultivation of shallot cv. Tajuk in Bantul and Kulon Progo areas. This study also collected information about cropping system applied in each area to determine its effect on the prevalence of Fusarium disease. The results of this study are expected to provide beneficial information regarding the resistance of Tajuk variety against Fusarium disease. This information would become a reference required to develop better recommendations for the cultivation of shallot cv. Tajuk, particularly in Bantul and Kulon Progo areas.

### **MATERIALS AND METHODS**

This study was conducted in the selected shallot fields in Bantul and Kulon Progo Regency. Selection of shallot fields used in this study was based on the use of shallot cv. Tajuk for the cultivation. Eight shallot fields in Bantul Regency and six shallot fields in Kulon Progo Regency were carefully selected to be used (Table 1) in disease monitoring and survey of cropping system. For Bantul Regency, disease observation and cropping system survey at each selected field were carried out during dry season from June to August 2023. For Kulon Progo Regency, both surveys were performed during rainy season from January to February 2024. Cropping system applied in each selected shallot field was also collected through land-owner interview. Several aspects in shallot cultivation were identified, such as cropping pattern, cropping frequency, use of crop rotation, plant spacing, and watering intensity.

Table 1. Details of shallot fields used in this study

Regency	District	Sub-village	Location Coordinate	Elevation* (m asl)
Bantul –	Imagiri	Kalidadap	7°57'51.2"S 110°22'48.9"E	113.64
	Imogiri	Srunggo	7°58'17.5"S 110°22'53.6"E	169.03
		Srigading 1	7°59'57.7"S 110°15'53.4"E	5.38
	Sanden	Gadingsari	8°00'02.5"S 110°15'33.3"E	7.89
		Srigading 3	7°59'18.2"S 110°16'32.1"E	7.27
	Kretek	Kretek 1	7°59'42.3"S 110°18'54.2"E	10.42
		Kretek 2	7°59'39.8"S 110°18'47.8"E	8.65
		Sono	7°59'54.5"S 110°18'41.6"E	7.36
Kulon Progo		Demangrejo	7°52'24.4"S 110°12'25.1"E	25.03
	Sentolo	Karasan	7°52'30.5"S 110°12'13.8"E	25.48
		Kalimenur	7°52'10.2"S 110°12'31.7"E	26.63
	Lendah	Ngentakrejo 1	7°53'45.7"S 110°14'58.3"E	22.79
		Ngentakrejo 2	7°53'31.0"S 110°15'07.8"E	25.35
		Ngentakrejo 3	7°53'49.7"S 110°15'12.2"E	24.69

Remarks: \*Elevation was identified using Google Earth.

Monitoring of Fusarium disease occurrence was performed using protocol adapted from Aisyah et al. (2022) with slight modification. Disease symptom was observed weekly starting from the plants aged 4 weeks until 7 weeks. About 20 plants collected from 5 different points in each field were collected to observe the disease based on the scoring of leaf symptoms (Table 2). Severity of Fusarium disease was further determined using the following formula [1]:

Disease Severity (%) = 
$$\frac{\sum_{i=0}^{n}(n_i \times v_i)}{N \times V} \times 100$$
 ......[1]

where  $n_i$  is the number of symptomatic plants representing each score,  $v_i$  is the score of severity, N is the number of plants observed, and V is the highest symptom score. Data obtained from disease monitoring was statistically analyzed using one-way analysis of variance. Statistical analysis was conducted using SPSS version 26.0. Significance recorded among location was subsequently analyzed using Duncan's New Multiple Range Test (DNMRT) with a p<0.05.

**Table 2.** Scoring used to measure leaf twisting symptom on infected shallot

Score	Symptom Description			
1	No visible symptom			
3	Chlorosis of the youngest leaves			
5	Necrosis on the leaf tip			
7	Elongated (excessive length increase) leaf followed with yellowing			
9	Leaves dry out and twisted			
12	Plant dead			

Source: Herlina et al. 2021

### RESULTS AND DISCUSSIONS Results

Shallot cv. Tajuk grown in Bantul and Kulon Progo showed different responses towards the infection of *Fusarium* disease indicated by the difference of disease development dynamics (Figure 1). Development of *Fusarium* disease in Bantul fields was more progressive than those recorded in Kulon Progo. Exponential increase of disease severity at 5 WAP was shown in Imogiri, while Sanden revealed higher severity since 4 WAP (Figure 1). However, the severity of *Fusarium* disease in this area became stagnant at 6 and 7 WAP. Contrast pattern was shown in Kulon Progo fields where the disease slowly developed from 4 to 7 WAP (Figure 1) resulting in less severe impact.


Severity of *Fusarium* disease at 7 WAP from each shallot field in Bantul and Kulon Progo was compared to determine the diversity in resistance of this variety within the same site. Of all fields observed in Bantul, the most severe infection was found in Kretek 2, exhibiting a significant difference compared to other villages, such as Kretek 1, Srigading 1, Srigading 3, and Srunggo (Figure


1). Interestingly, disparity of disease severity among shallot fields in Kretek District was more remarkable than those recorded in Sanden and Imogiri. Such notable gap in disease severity within the same district was also found in Kulon Progo. Figure 2 demonstrated significant difference in severity between Kalimenur and Ngentakrejo 1 compared to other fields within the same district. Shallot cv. Tajuk cultivated in Ngentakrejo 1 showed the lowest disease severity (13.3%), while those cultivated in Kalimenur exhibited the highest severity of about 25% (Figure 2).

Occurrence of plant disease is affected by various factors, including cultural practices applied. Table 3 and **Table 4** demonstrated the difference of cultural practices applied during the cultivation of shallot cv. Tajuk in Bantul and Kulon Progo. Most shallot fields in both regencies applied intercropping with chili, while all observed fields in Bantul and Kulon Progo used rotation (Table 3 and Table 4). In terms of planting frequency, shallot cultivation in Bantul varied among districts. District in Bantul with the most frequent shallot planting was Sanden (particularly Srigading 1 and Gadingsari), where fields observed in Imogiri cultivated shallot once in a year only (Table 3). Unlike Bantul, all fields observed in Kulon Progo cultivated shallot twice a year (Table 4).


Other aspects of farming practices observed in this study were plant spacing and watering intensity. Plant spacing used for shallot planting varies among districts in both areas. Several districts applied the same plant spacing in all observed fields, as shown in Imogiri and Kretek (Bantul) (Table 3), as well as Lendah (Kulon Progo) (Table 4). Distinct variation in plant spacing among shallot fields was found in Sanden (Table 3). In terms of watering frequency, most shallot fields in Bantul performed watering every two days, except in Srigading 3 (Table). In contrast, all shallot fields observed in Kulon Progo applied daily watering (Table 4). Referring to the observation period for each area, the watering frequency implemented in both areas appeared to contradict the weather conditions, particularly rainfall. Observation in Bantul was carried out during the dry season (June -August 2023), where the rain intensity within this period was extremely low (Table 5). On the other hand, the rainfall recorded in the Kulon Progo area during the observation period (January – February 2024) was high (Table 5).


Based on the disease severity at 7 WAP, the most severe *Fusarium* infection in Bantul was recorded from shallot field in Kretek 2 (Figure 2). Shallot cv. Tajuk cultivated in this village applied intercropping with chili, quite wide plant spacing, and used less frequent watering (Table 3) during the observation period. The highest severity of *Fusarium* disease from shallot fields in Kulon Progo was found in Kalimenur, while the lowest severity was recorded in Ngentakrejo 1 (Figure 2). Cultivation of shallot cv. Tajuk in Kalimenur utilized narrower plant spacing (15x15 cm) and implemented rotation with paddy and chili (Table 4). In contrast, the shallot field in Ngentakrejo 1 applied wider plant spacing (20x18 cm) and performed rotation with chili and peanut (Table 4).





**Figure 1.** Comparison of *Fusarium* disease development between shallot cv. Tajuk fields in each district of Bantul (a) and Kulon Progo (b)





**Remarks:** Bars with the same letters shows insignificant difference based on DNMRT with a p<0.05.

Figure 2. Severity of Fusarium disease at 7 WAP in each shallot field representing Bantul (a) and Kulon Progo (b).

Table 3. Differences of shallot cv. Tajuk cultivation practices applied in several fields in Bantul Regency

	Aspects of farming practices					
District/Village	Cropping system	Intercropped crops	Planting frequency (times.year <sup>-1</sup> )	Rotated crops	Plant spacing (cm x cm)	Watering frequency
Imogiri District						
Kalidadap	Intercropping	Chili	- 1	Paddy, chili, tobacco	- 20 x 20	Once in 2 days
Srunggo	Monoculture	None		Paddy, tobacco		
Sanden District						
Srigading 1	Intercropping	Chili	3	Watercress, eggplant	20 x 20	Once in 2
Gadingsari					19 x 19	days
Srigading 3			2	Chili, paddy	15 x 20	Daily
Kretek District						
Kretek 1		Chili, peanut				
Kretek 2	Intercropping	Chili	2	Chili, paddy	20 x 20	Once in 2 days
Sono		Offilii				

Table 4. Differences of shallot cv. Tajuk cultivation practices applied in several fields in Kulon Progo Regency

	Aspects of farming practices						
District/Village	Cropping system	Intercropped crops	Planting frequency (times/year)	Rotated crops	Plant spacing (cm x cm)	Watering frequency	
Sentolo District							
Demangrejo 1	Monoculture	None	- 2	Chili	18 x 18	- Daily	
Karasan	- Intercropping	Chili			- 15 x 15		
Kalimenur				Paddy, chili			
Lendah District							
Ngentakrejo 1				Chili, peanut			
Ngentakrejo 2	Intercropping	Chili	2	Chili	20 x 18	Daily	
Ngentakrejo 3	-			Cilli			

Table 5. Monthly rainfall and number of rainy days in Bantul and Kulon Progo during the observation period

Regency	Observation Period	Average Rainfall (mm) per Month	Number of Rainy Days per Month	
		0.0 (June)	0.0 (June)	
Bantul	June – August 2023	24.0 (July)	5.0 (July)	
		0.0 (August)	1.0 (August)	
Kulan Drago	January Fahruary 2024	405.0 (January)	22.0 (January)	
Kulon Progo	January – February 2024	147.0 (February)	10.0 (February)	

Source: Statistics of Yogyakarta Province (2024a).

#### Discussions

Bantul and Kulon Progo are the main contributors to regional shallot production. Statistics of Yogyakarta Province (2024b) reported that shallot production in both regencies in 2023 reached 94% of total shallot production in the Special Region of Yogyakarta Province. However, shallot production in both Bantul and Kulon Progo has not reached its optimal productivity due to various issues during its cultivation, including disease infection. Fusarium disease is one of the important diseases infecting Allium species, including shallot (Le et al. 2021; Poursakhi et al. 2024). In Indonesia, disease caused by Fusarium spp. in shallot, garlic, and leeks is mainly known as moler or twisted disease since the symptom that appeared is leaf twisting.

Prevalence of *Fusarium* disease in shallots may vary depending on variety, environmental factors, and pathogen species/strain. This present study highlighted the *Fusarium* intensity recorded from the shallot cv. Tajuk cultivated in several areas in Bantul and Kulon Progo, where the disease observation was carried out at different times between Bantul and Kulon Progo. Shallot cv. Tajuk cultivated in Bantul conferred higher susceptibility to *Fusarium* disease than those cultivated in Kulon Progo (Figure 2). In addition, Figure 1 also emphasized the distinct differences in disease progression from 4 to 7 WAP between Kulon Progo and Bantul. These findings suggest that shallot fields in Bantul may experience higher exposure to this disease than those in Kulon Progo.

Regarding the disease severity, the highest severity (43.8%) in Bantul was recorded from shallot field in Kretek 2 (Figure 2). This finding was higher than those reported by Wibowo et al. (2023), where the twisted disease infecting shallot cv. Tajuk cultivated in the coastal area of Bantul reached its highest incidence at 33.97% during the rainy season in 2021. Interestingly, this present study found that high disease severity in Bantul occurred during the dry season with extremely low rainfall (Table 5). Moreover, result of this study implies that the increase in disease severity recorded in Bantul compared to the previous study might be associated with the level of pathogen infestation in the specified field. The longer the pathogen is infested and survives in the field, the higher the chance of disease severity increases in the next infection cycle. Despite the accumulation of pathogen residue, another factor that contributes to more severe infection is the possibility of complex Fusarium species. Infection by these complex species may result in stronger pathogenicity and higher virulence, leading to a more severe impact on plant growth (Sidharthan et al. 2023). Other studies also highlighted that the composition of Fusarium complex varied by location and host, hence resulting in different levels of virulence (Degani et al. 2024; Le et al. 2021).

Considering the shallot variety, resistance of Tajuk variety to *Fusarium* infection has been investigated previously under greenhouse conditions (Hadiwiyono et al. 2020; Herlina et al. 2021). This variety was considered to be in the moderate category for infection

by Fusarium oxysporum f.sp. cepae resulting in 37.17% severity (Hadiwiyono et al. 2020). In contrast, Herlina et al. (2021) reported higher severity (68.54%) from this shallot variety after being inoculated with three isolates of FOC. Comparison of these previous studies underlines that the susceptibility of the shallot variety to Fusarium disease could fluctuate depending on Fusarium species. In that context, Fusarium has been known for its diverse host range due to the formation of formae speciales (f.sp.) with different virulence and host specificity (Edel-Hermann and Lecomte 2019; Gordon 2017). However, the response of Tajuk variety to specified Fusarium species/strains may differ depending on geographical location since various environmental aspects may affect disease development. Therefore, further research on the resistance of this variety across multiple locations in Indonesia would provide an in-depth knowledge regarding the characteristics of Tajuk variety.

Findings of this study conveyed the variation of cultural practices applied in the cultivation of Tajuk variety between Bantul and Kulon Progo (Table 3 and Table 4). Most of the representative shallot fields in each regency applied an intercropping system with chili as the intercropped plant. Our previous work in 2021 also found that the cultivation of this shallot variety in Kretek and Sanden (Bantul Regency) was performed with the same choice of intercropped crop (Aisyah et al. 2022). Despite the intercropping system applied in most shallot fields in this study, the occurrence of Fusarium disease showed higher severity in Bantul Regency than Kulon Progo (Figure 1 and Figure 2). It implies that the use of intercropping and crop rotation in Bantul is unable to mitigate the disease prevalence significantly. Such high disease severity recorded from most shallot fields in Bantul indicates that the disease prevalence in this area has reached the state of endemic. Therefore, regardless of the variation of farming practices applied, the disease severity did not differ significantly. In addition, this endemic state was also marked with high severity developed during the dry season in Bantul (Table 5) and low watering intensity applied during the shallot cultivation (Tabel 3a).

Endemic of Fusarium disease in a specific geographical area could occur due to the establishment of the pathogen during several growing seasons. This establishment could result in higher virulence depending on the diversity of species or the presence of a species complex (Delgado-Ortiz et al. 2016; Haapalainen et al. 2016). Infection by these complex species has a higher chance to cause greater growth reduction and yield loss (Sidharthan et al. 2023). Interestingly, complex species of Fusarium may consist of both pathogenic and nonpathogenic species and the formation of this species complex can be triggered through crop rotation. Diabankana et al. (2024) reported the infection of Fusarium strains isolated from onion on other plant species used for crop rotation, suggesting the risk of Fusarium co-infection to non-host plants. Aligned with this previous study, all shallot fields in Bantul applied rotation with various non-Allium crops (Table 3) with no significant impact on the disease severity (Figure 2). Nonetheless, the Fusarium species infecting each shallot field in Bantul and Kulon Progo has not been thoroughly

explored in this present study. Further investigation is recommended to identify whether the *Fusarium* disease found in both locations is caused by a single or complex species.

Contrast disease prevalence recorded from shallot fields in Kulon Progo, highlighting the possibility of lower exposure of *Fusarium* in this area, thus resulting in less severe impact on shallot cv. Tajuk (Figure 1 and Figure 2). Considering the cultural practices applied, shallots cultivated using both monoculture and intercropping systems showed similar disease severity in this region (Figure 2). Nevertheless, the lowest disease severity (13.3%) in Kulon Progo was obtained from Ngentakrejo 1 (Lendah District) (Figure 2) where the cultivation applied intercropping with chili and used peanut as one of the rotated crops (Table 4). This finding is higher compared to those reported by Pustika et al. (2024), where Tajuk variety cultivated in Panjatan District (Kulon Progo) showed 7.60% severity (at 8 WAP) to Fusarium infection recorded in 2022. Not only the location difference, but also the difference in Tajuk variety resistance to *Fusarium* disease between this present study and the previous one could be attributed to the climatic conditions during the observation. Disease monitoring conducted by Pustika et al. (2024) was carried out during the dry season (August to October 2022), while the present study was observed during the rainy season (January to February 2024). However, to justify the prevalence of this Fusarium disease in a specified area more accurately, disease monitoring should be conducted in a specified field continuously following the growing season of shallot in that area. Thus, the dynamic of the disease incidence in each growing season could be comprehensively quantified to determine the factors affecting its severity as well as the resistance of the shallot variety.

Overall, the findings of this study highlight the susceptibility of shallot cv. Tajuk to Fusarium disease varies depending on the location of its cultivation, instead of the cultural practices applied in both Bantul and Kulon Progo. The difference in disease severity recorded in each location suggests the different levels of disease exposure among locations that could be linked to the pathogen establishment and virulence. Further analysis should focus on the identification of Fusarium species infecting each shallot field in both locations to evaluate the spectrum of pathogen virulence. In addition, this present study also reveals that higher rainfall does not appear to be in line with the increased disease intensity, thus indicating a shift in the interaction of plant-pathogenenvironment due to climate change effects. Nonetheless, deeper investigation is needed to complement the findings of this study through consecutive disease monitoring following the shallot cv. Tajuk growing season in both locations.

# CONCLUSIONS AND SUGGESTIONS Conclusions

Balok Merah accession showed the best results, supported by the character of the highest maximum growth potential, germination power, and dry weight of normal sprouts, and more consistent moisture content. The storage method with panicles showed the best

results in maintaining the quality of seeds during storage, supported by the character of the highest maximum growth potential and germination power. Proximate analysis showed that the highest moisture content, ash content and protein content were found in the Balok Merah. The highest fat content in the accession of Pulut Hitam Rice. Highest carbohydrate levels at Mukot accession. There is a strong positive correlation between maximum growth potential and germination power.

#### **Suggestions**

West Bangka local rice accession in the form of Balok Merah can be stored for 8 months. The use of effective storage methods, such as panicles or airtight packaging (cans), can be applied by farmers and the public in storing Bangka local rice accession seeds to maintain seed quality. Further research is needed to explore other storage methods with various local rice accessions in Bangka to support agricultural productivity in Bangka Belitung.

### **REFERENCES**

- Aisyah SN, Khoiruddin M, Hidayat T, Astuti A. 2022. Comparison on Fusarium basal rot occurrence among shallot cultivations in Bantul Regency. IOP Conf Ser Earth Environ Sci. 985:012052. https://doi.org/10.1088/1755-1315/985/1/012052.
- Anggrasari H, Dhamira A, Maheresmi H. 2025. Shallot price forecast in traditional markets of the Special Region of Yogyakarta. SHS Web Conf. 212:04013. https://doi.org/10.1051/shsconf/202521204013.
- Arifin Z, Gunawan I, Nursamukti A. 2025. Enhancing shallot cultivation efficiency through smart farming: Drone technology for precision land mapping and yield estimation. IOP Conf Ser Earth Environ Sci. 1490:012034. https://doi.org/10.1088/1755-1315/1490/1/012034.
- Degani O, Dimant E, Margalit E. 2024. Impact of fusarium species composition and incidence on onion basal rot in northeastern israel. Horticulturae. 10(4):373. https://doi.org/10.3390/horticulturae10040373.
- Delgado-Ortiz JC, Ochoa-Fuentes YM, Cerna-Chávez E, Beltrán-Beache M, Rodríguez-Guerra R, Aguirre-Uribe LA, Vázquez-Martínez O. 2016. Patogenicidad de especies de Fusarium asociadas a la pudrición basal del ajo en el Centro Norte de México. Rev Microbiol. 48(3):222-228. https://doi.org/10.1016/j.ram.2016.04.003.
- Dewi FS, Dewi RR, Abadi AL, Setiawan A, Aini LQ, Syib'li MA. 2025. Biocontrol of Fusarium oxysporum f.sp. cepae on Indonesian local garlic plants (Lumbu Hijau) using a consortium of Bacillus amyloliquefaciens B1 and arbuscular mycorrhizal fungi. Mycobiology.1-9. https://doi.org/10.1080/12298093.2024.2433826.
- Diabankana RGC, Frolov M, Islamov B, Shulga E, Filimonova MN, Afordoanyi DM, Validov S. 2024. Identification and aggressiveness of Fusarium species associated with onion bulb (Allium cepa L.)

- durina Funai. 10(2):161. storage. https://doi.org/10.3390/jof10020161.
- Dinata GF, Ariani N, Purnomo A, Aini LQ. 2021. Pemanfaatan biodiversitas bakteri serasah kopi sebagai solusi pengendali penyakit moler pada bawang merah. J HPT. 9(1):28-34. http://dx.doi.org/10.21776/ub.jurnalhpt.2021.009.1.5.
- Edel-Hermann V, Lecomte C. 2019. Current status of Fusarium oxysporum formae speciales and races. Phytopathology. 109(4):512-530. https://doi.org/10.1094/phyto-08-18-0320-rvw.
- FAOSTAT. 2023. Onion production, productivity. http://www.fao.org/faostat/en/#data/QC.
- Fauzia W, Maryani Y, Darnawi D. 2020. The effect of various kinds organic fertilizers on growth and yield of shallot (Allium ascalonicum L.) Thailand and sarmo varieties. Jurnal Ilmiah Agroust. 4(1):66-75.
- Gordon TR. 2017. Fusarium oxysporum and the fusarium wilt syndrome. Annu Rev Phytopathol. 55(1):23-39. https://doi.org/10.1146/annurev-phyto-080615-095919.
- Gusman TA, Zakiyyah Z, Nurudin A, Yulina IK. 2025. Development herbal syrup from shallot skin: The effect of cmc addition for bioactive compounds and proximate content. E3S Web Conf. 622:02007. https://doi.org/10.1051/e3sconf/202562202007.
- Haapalainen M, Kuivainen E, Iivonen S, Niemi M, Latvala S. 2023. Pathogenicity of Fusarium oxysporum and Fusarium proliferatum isolates from symptomless onions (Allium cepa) and onions with Fusarium basal rot. Plant Pathol. 72(6):1122-1135. http://dx.doi.org/10.1111/ppa.13718.
- Haapalainen M, Latvala S, Kuivainen E, Qiu Y, Segerstedt M, Hannukkala A. 2016. Fusarium oxysporum, F. proliferatum and F. redolens associated with basal rot of onion in Finland. Plant 65(8):1310-1320. http://dx.doi.org/10.1111/ppa.12521.
- Hadiwiyono, Sari K, Poromarto SH. 2020. Yields losses caused by basal plate rot (Fusarium oxysporum f. sp. cepae) in some shallot varieties. Caraka Tani. 35(2):250-257. https://doi.org/10.20961/carakatani.v35i2.26916.
- Herlina L, Istiaji B, Wiyono S. 2021. The causal agent of Fusarium disease infested shallots in Java Islands of E3S Web Indonesia. Conf. 232:03003. https://doi.org/10.1051/e3sconf/202123203003.
- Le D, Audenaert K, Haesaert G. 2021. Fusarium basal rot: Profile of an increasingly important disease in Allium spp. Trop Plant Pathol. 46:241–253. https://doi.org/10.1007/s40858-021-00421-9.
- Mahanty B, Mishra R, Joshi RK. 2023. Fusarium oxysporum f. sp. cepae small RNAs (Foc-sRNAs) promote disease susceptibility in onion (Allium cepa L.) through cross kingdom RNA interference. Physiol 125:102018. Mol Plant Pathol. https://doi.org/10.1016/j.pmpp.2023.102018.

- Mariani M, Suprapta DN, Sudana IM, Temaja IGRM, Sudantha IM. 2022. First report of *Nectria haematococca* causing a moler disease on shallots in West Nusa Tenggara, Indonesia. Biodiversitas. 23(5). http://dx.doi.org/10.13057/biodiv/d230559.
- Neda N, Zahra BH, Mohammad RH, Mitra A, Mohammad RM, Akram GA, Maryam M-S, Mehdi M. 2025. Promising effects of Persian shallot extract on the serum markers and blood pressure of patients with metabolic syndrome: A double-blinded randomized controlled trial. J Tradit Chin Med. 45(1):100. https://doi.org/10.19852/j.cnki.jtcm.2025.01.008.
- Pasaribu S. 2017. Pengaruh dosis pupuk kompos gulma siam terhadap pertumbuhan dan hasil tiga varietas bawang merah [undergraduate thesis]. Yogyakarta (ID): Universitas Mercu Buana.
- Poursakhi S, Asadi-Gharneh HA, Nasr-Esfahani M, Abbasi Z, Khankahdani HH. 2024. Identification of novel associations of candidate marker genes with resistance to onion-*Fusarium* basal rot interaction pathosystem. Plant Gene. 37:100440. https://doi.org/10.1016/j.plgene.2023.100440.
- Pustika A, Purwaningsih H, Rahmawati R, Darini M. 2024. Fusarium wilt of some shallot local varieties in yogyakarta alternating-bed (surjan) agroecosystem. IOP Conf Ser Earth Environ Sci. 1377:012108. https://doi.org/10.1088/1755-1315/1377/1/012108.
- Ridqi MF, Kusumaningsih A. 2025. Income risk of shallot farming in Selopamioro Village, Imogiri Sub-district, bantul Regency. IOP Conf Ser Earth Environ Sci. 1518:012023. http://dx.doi.org/10.1088/1755-1315/1518/1/012023.
- Sharma S, Mandal S, Cramer CS. 2024. Recent advances in understanding and controlling *Fusarium* diseases of *Alliums*. Horticulturae. 10(5):527. https://doi.org/10.3390/horticulturae10050527.
- Sholeh MI, Nurcahyanti SD. 2023. Development of moler disease (*Fusarium Oxysporum* F.Sp *Cepae*) at The Shallot Production Center in Probolinggo Regency. 6(2):56–62. http://dx.doi.org/10.19184/bip.v6i2.35392.
- Sidharthan VK, Pothi-Raj G, Suryaprakash V, Singh AK, Aggarwal R, Shanmugam V. 2023. A synergic and compatible microbial-based consortium for biocontrol of *Fusarium* wilt of tomato. Phytopathol Mediterr. 62:183-197. http://dx.doi.org/10.36253/phyto-13055.
- Sogoba KH, Kadidia K, Alassane O, Mohamed S, Itolou KA, Harouna S, Claudine C. 2022. Pathogenic characterization of three *Fusarium* species associated with onion (*Allium cepa* L.) in Burkina Faso.
  - http://dx.doi.org/10.33687/phytopath.011.03.4327.
- Statistics Indonesia. 2025. Agricultural statistic for horticulture: Production of annual vegetable and fruit crops in Yogyakarta Province 2024. June 10th, 2025.
- Statistics of Yogyakarta Province. 2024a. Curah hujan per bulan (mm), 2023.

- Statistics of Yogyakarta Province. 2024b. Statistics of horticulture in Yogyakarta Province 2023.
- Sutardi, Kristamtini, Purwaningsih H, Widyayanti S, Arianti FD, Pertiwi MD, Triastono J, Praptana RH, Malik A, Cempaka IG. 2022. Nutrient management of shallot farming in sandy loam soil in Tegalrejo, Gunungkidul, Indonesia. Sustainability. 14(19):11862. https://doi.org/10.3390/su141911862.
- Wibowo A, Santika IA, Syafitri LM, Widiastuti A, Subandiyah S, Harper S. 2023. Incidence of twisted disease and cultivation practice of shallot farmers in Bantul Coastal Area, Yogyakarta, Indonesia. J Trop Plant Pest Dis. 23(1):23-30. https://doi.org/10.23960/jhptt.12323-30.