

Control of Bacterial Leaf Blight *Xanthomonas* by Application of *Bacillus subtilis* Biobactericide on Rices

Iqbal Erdiansyah*, Khusnul Khotimah

Food crop Production Technology Study Program, Department of Agricultural Production, Politeknik Negeri Jember, Jember, East Java 68121, Indonesia

Received August 27, 2025 Accepted September 28, 2025

ABSTRACT

Indonesia's agricultural problems in rice cultivation are increasingly complex, threatening national food security. The attack of *Xanthomonas oryzae* pv. *oryzae* (Xoo) can cause decreased production of bacteria, resulting in rice disease bacterial leaf blight (BLB). The biological agent *Bacillus subtilis* (*B. Subtilis*) has the potential to control BLB disease. The study aimed to determine the effect of *B. subtilis* bacteria in reducing the intensity of Xoo attacks by comparing the use of pesticides. Xoo attacks by comparing the use of citronella vegetable pesticide and its effect on rice production. This experiment was designed as a comparison of 2 plots with different treatments, namely the control treatment of citronella vegetable pesticide with a concentration of 75% and the biobactericide treatment of *B. subtilis* with an antagonistic bacterial suspension density of 33×10^4 cfu ml⁻¹. The variables observed were inhibition, disease intensity, weight of harvested dry grain (HDG) per clump, and weight of HDG per plant. The results stated that the biobactericide *B. subtilis* had a real effect on the development of BLB disease up to an intensity of 14.72% and the development of BLB with citronella vegetable pesticide treatment decreased to 28.97%, the weight of HDG per clump and did not have a real effect on the weight of HDG per panicle.

Keywords: Antagonistic bacteria; Biobactericide, Citronella vegetable pesticide; Disease intensity; Harvested dry grain; Leaf blight

Cite this as (CSE Style): Erdiansyah I, Khotimah K. 2025. Control of bacterial leaf blight *Xanthomonas* by application of *Bacillus subtilis* biobactericide on rices. Agrotechnology Res J. 9(2):90–95. <https://dx.doi.org/10.20961/agrotechresj.v9i2.91028>.

INTRODUCTION

Rice (*Oryza sativa* L.) is the main food in Indonesia, because the majority of the population uses rice as their staple food (Donggulo et al. 2017). According to data from BPS Jember (2023), shows that rice production in Jember Regency from January to December 2022 reached approximately 607.37 thousand tons of dry paddy (GKG), with a predicted decrease of approximately 8.33 thousand tons of GKG, or 1.35%, compared to 615.70 thousand tons of GKG in 2021. With the high level of Indonesian people's reliance on rice as a staple food, the community may eventually shift to a non-rice diet, so the government needs to achieve rice self-sufficiency (Moningka et al. 2020). The complexity of the problems faced by the agricultural industry, especially rice crops, places this country's food security at the forefront of its problems.

Examples of factors that threaten food security include agricultural land conversion, inappropriate use of chemical pesticides, labor force decline, climate change,

and pest attacks (Molotoks et al. 2021; Wakil et al. 2025). One important disease affecting rice plants is leaf blight (LBB), caused by the bacterium *Xanthomonas oryzae* pv. *oryzae* (Xoo). Loss of rice production due to bacterial blight disease is around 15-24% (Sutarman 2017). Plant symptoms include yellowing, drying, and straw-colored leaves, commonly known as leaf spots. Several disease control methods have been implemented, including land sanitation, the use of resistant cultivars, synthetic chemical pesticides, biopesticides, and biological agents.

The use of chemical pesticides can be reduced by using biopesticides made from natural and environmentally friendly ingredients. Plants such as citronella (*Cymbopogon nardus*) can act as natural antimicrobials. The use of biological agents, such as those from the *Bacillus subtilis* (*B. subtilis*) group, can also be an alternative solution, which has been tested in vitro to suppress the development of Xoo in rice seeds (Djatzmiko et al. 2022; Srinivas et al. 2024). Many scientists have reported how biological agents prevent disease and increase production.

Increasing production and preventing diseases in rice plants using biological agents of the *B. subtilis* group has not been widely reported, especially in Indonesia (Agustiansyah et al. 2013). Based on the explanation

*Corresponding Author:
E-Mail: iqbal@polije.ac.id

above, it is known that *B. subtilis* bacteria can act as a biological agent. If the antagonist produced by *B. subtilis* is high, Xoo disease attacks can be controlled. Therefore, research on *B. subtilis* bacteria was conducted to determine the effect of the biotoxin *B. subtilis* in controlling the intensity of Xoo disease in rice plants.

MATERIALS AND METHODS

This research was conducted in rice fields in Tlogo Wetan Village, Antirogo, Summersari District, Jember Regency, East Java and the Bioscience Laboratory, Jember State Polytechnic in August-December 2023. The tools and materials used in this research are, the tools used in the first stage of research include air flow cabinet (LAFC), sterile petri dishes, loop needles, laminar autoclaves, bunsen, test tubes, cell phone cameras, sprayers, measuring cups, hot plates, pH meters, thermoshakers, vortex, incubators, micropipettes, Eppendorf tubes, vernier calipers. While the tools used in the second stage of research include, sample boards, hoes, banners, meters, scales, measuring cups, and sprayers, cameras. The first phase of the research used materials such as *Bacillus* sp. bacterial isolate, Xoo bacterial isolate, 96% alcohol, 70% alcohol, fine rice bran, distilled water, aluminum foil, chloroform, filter paper (Whatman paper no. 1), sterile tissue, label paper, plastic, camera. In the second phase, among others, were media resulting from the multiplication of *B. subtilis* bacteria, Xoo bacterial isolate, topsoil, Ciherang variety rice seeds, manure, botanical pesticides, NPK fertilizers and sample labels. The research design compared two treatment plots on different rice fields, where each field was sprayed with a biological agent and botanical pesticide. Plot 1 will be sprayed using botanical pesticides used when controlling diseases on organic fields with a concentration of 5 ml of citronella leaf extract plus 5 ml of water (Barimbing 2022). Plot 2 was sprayed with the biological agent *B. subtilis* at a concentration of 20 mL.L⁻¹. The field research layout was 10 m x 10 m for the plot and 25 cm x 25 cm for the planting distance, with 2-3 seedlings per hole (Erdiansyah et al. 2021). The random sampling method used was a zigzag pattern with a sample of 50 plants.

Bacillus subtilis propagation in the laboratory uses rice bran as a medium. The mechanism for making rice bran media is to weigh 30 grams of processed, fine rice bran and sterilize it using an autoclave for 30 minutes at a temperature of 121°C and a pressure of 1 atm.

Antagonistic tests were conducted between *B. subtilis* bacteria and the pathogenic bacteria that cause bacterial leaf blight, namely Xoo, to determine the potential of pure bacteria as biological agents. The method used was the modified Kawaguchi (fogging) method in NA media. Adding 1 mL.100 mL of starter and incubating for 3 days, then it was ready to be applied. The media resulting from the incubated bacterial multiplication was applied to the plants. Application to rice seedlings was carried out when they were about to be transplanted, by dipping them in a *B. subtilis* biobactericide solution for 10 minutes and ready to be planted.

Spraying is carried out on the seedbed (seedbed) approximately 7 days before transplanting.

Biobactericide spraying is carried out after planting at 2, 3, 4, and 5 weeks after planting. The data obtained were analyzed using non-parametric statistical tests in SPSS with the following stages: 1) Data normality test; 2) Data homogeneity test; 3) If the data is not normal and not homogeneous, then it is continued with a non-parametric test using the Mann-Whitney Test; 4) If the data is normal and homogeneous, then it is continued with the Paired-Samples T Test.

RESULTS AND DISCUSSIONS

Controlling disease with chemical bactericides is very harmful to the environment. Natural bactericides are needed as an alternative to minimize the harmful impact on the environment. Biobactericides are substances or components that can inhibit bacterial growth. Soil bacteria called *Bacillus* spp. are often found in the rhizosphere of plants (Pandango et al. 2018). Because *Bacillus* rhizobacteria can control plant diseases and increase the amount of nutrients in the soil, its use as a biological agent has been shown to increase yields (Nurkartika et al. 2017).

One bacterial capable of addressing root problems is *Bacillus*. This bacterium is safe to use as a biobactericidal agent and grows rapidly even in liquid media. As a plant growth-promoting rhizobacterium (PGPR) and able to survive harsh climatic conditions, *B. subtilis* has an advantage over other similar bacteria, namely its ability to reduce various plant diseases (Wartono and Mutaqin 2015; Serrão et al. 2024; Ali et al. 2025). Based on the results of the research data, the following results were obtained.

Inhibitory power

To determine the potential of *B. subtilis* bacteria, an inhibitory power test was conducted using the spray method. The results were measured in millimeters (mm). The inhibitory power test results are presented in Figure 1. The pathogenic bacteria Xoo was suppressed in its development by the antagonistic bacteria *B. subtilis*, based on the findings of an antagonist test conducted in vitro using NA media in a petri dish. An indication of the ability of the pathogen Xoo to be inhibited by a biological agent is the formation of an inhibition zone around the antagonist colony. The width of the inhibition zone, or bacterial inhibition zone, indicates how well the microorganism can inhibit the growth of the disease (Azman et al. 2017; Srinivas et al. 2024).

The antibacterial effect is divided into four levels: weak, moderate, strong, and very strong. If the inhibition zone diameter is <5 mm, bacterial activity is considered weak, moderate if 5-10 mm, strong if 10-20 mm, and very strong if more than 20 mm (Barimbing 2022). Based on the results of measuring the diameter of the clear zone in the *B. subtilis* bacterial isolate against the Xoo bacteria. Both replications had an inhibitory power in the moderate inhibition category, namely 5.5 mm in the first replication and 5.5 mm in the second replication. Based on the antagonist test, it can be seen that the *B. subtilis* isolate can inhibit the development of the Xoo bacteria. The antagonist bacteria *B. subtilis* has an inhibitory substance so that the growth of the antagonist pathogenic bacteria can be inhibited.




Figure 1. (a) Inhibitory power test of *B. subtilis* bacteria and *Xoo* bacteria, (b) Results of inhibitory power test of *B. subtilis* bacteria and *Xoo* bacteria

The resulting inhibitory compounds act as pathogen antagonists and work by degrading the cell walls of pathogenic bacteria. Previous research has shown that citronella contains essential oils with potent antibacterial properties. Citronella has an inhibition zone of 10.00 mm at a 50% concentration, which is classified as moderate inhibition (Barimbing 2022). The primary active compounds in citronella essential oil include citronellal, citral, geraniol, and myrcene that exhibit natural antimicrobial properties.

Table 1. Results of inhibitory power test calculations based on the method of

Resistance (mm)	Diameter	Inhibition Zone Category
5.5 mm		Strong Category

Remarks: Weak category (≤ 5 mm), moderate category (6-10 mm), strong category (19-20 mm), and very strong category (≥ 20 mm).

Disease intensity

Observations showed that inoculation of *Xoo* isolates can cause symptoms on leaves 3 weeks after inoculation. Environmental factors such as temperature can influence the development of diseases such as bacterial leaf blight. The average daily temperature in the area where this experiment was conducted ranged from 23°C to 33°C, which supports the development of *Xoo* bacteria in rice plants because 26-30 °C is the ideal temperature for the development of these bacteria.

Based on the data in Table 2, the results of the intensity of bacterial leaf blight disease attacks (*Xoo*) between the treatment of the biological agent *B. subtilis* and the control treatment of the citronella leaf botanical pesticide provided significant results. The results of the disease intensity in the *B. subtilis* biobactericide treatment were lower compared to the disease intensity with the citronella botanical pesticide treatment. This can occur because the density of the formulation can suppress the rate of infection according to (Wartono and Mutaqin 2015), the reduction in bacterial leaf blight disease is influenced by the density of the formulation spraying. The density of the antagonistic bacterial suspension used was 33x cfu.ml⁻¹. The level of bacterial leaf blight disease intensity is in the moderate category because it is below 25%.

Table 2. Intensity of bacterial leaf blight disease (*Xoo*)

Treatment	Weeks After Planting (WAP)		
	3	4	5
Citronella Botanical Pesticide	15.22a	20.22a	28.97a
<i>B. subtilis</i>	16.73a	19.68b	14.72b

Remarks: Numbers followed by different letters in the same column indicate a significant difference according to the Paired samples T test ($P < 0.05$). WAP: weeks after planting

According to the Kementan (2018) the disease intensity levels follow the following provisions: (1) 0% No attacks/healthy plants, (2) 1-11% mild attacks, (3) 12-25% moderate attacks, (4) 26-85% severe attacks, and (5) >85% puso attacks. Therefore, control of bacterial leaf blight disease must be controlled. The effectiveness of these two bactericides is influenced by the active ingredients of the bactericide. This is in accordance with the statement Meimaharani and Listyorini (2013) that the use of *B. subtilis* plays a very important role in suppressing the population and intensity of disease attacks due to damage to plants. Botanical pesticides contain active ingredients of essential oils as antibacterials which can disrupt the process of causing pathogenic bacteria in leaves (Bota et al. 2015).

Observations were conducted before and after biobactericide application. Observations before application were conducted at 3 WAP with the aim of determining the rate of disease infection within the control threshold after inoculation of the pathogen *Xoo*. The decrease in the intensity of bacterial leaf blight disease is suspected to be due to the influence of compounds and toxins produced by the bacteria *B. subtilis* that infect bacterial leaf blight. At 4 WAP, the disease population increased again. Plant age influences the development of bacterial leaf blight disease; seedlings transplanted at a younger age often have a higher percentage of disease (Khaeruni et al. 2014).

Based on observations of disease severity, bacterial leaf blight disease progression in each treatment was determined on sample leaves. The disease severity showed that the earlier the rice plant was infected, the slower the disease spread; conversely, the earlier the rice plant was infected, the faster the disease spread. A decrease in disease severity was seen in the percentage of bacterial leaf blight disease at different periods after biobactericide application. Compared to the previous growth phase, disease progression occurred more rapidly during the vegetative growth phase, which lasted between 3 and 4 weeks after planting. The average disease progression reached 18-33%. The level of resistance of plant varieties is also influenced by the morphological structure of the leaf surface (Khaeruni et al. 2014). The slow rate of development in the reproductive phase is likely due to the complete formation of the plant's resistance structure. Plant epidermal cells have a waxy layer and a thick cuticle that are ideal for increasing plant resistance to diseases that directly penetrate the epidermal layer.

Dry rice harvest results

Test results showed that the application of the biobactericide *B. subtilis* significantly affected the weight of dry grain per clump in rice plants. The test results are presented in Table 3.

Table 3. Weight of dry paddy (GKS)

Treatment	Weight of Dry Paddy per Clump (g)
Citronella Botanical Pesticide	24.44a
<i>B. subtilis</i>	27.26b

Remarks: Numbers followed by different letters in the same column indicate significant differences according to the Mann-Whitney test ($P < 0.05$).

Table 3 shows that the *B. subtilis* treatment significantly affected the grain weight per hill. The results of the *B. subtilis* treatment were higher than those of plants treated with the citronella leaf pesticide, reaching an average weight of 27.26 grams. The percentage of dry paddy weight in the *B. subtilis* biobactericide treatment was 11.54% higher than that in the citronella plant pesticide treatment. This is due to the lower disease intensity in the biobactericide treatment. Production results and disease severity correlated consistently. Based on the research findings, population influences yield within a certain area.

The severity of the disease attack also affects the weight per sample; the greater the intensity of the attack caused by bacterial leaf blight, the greater the risk of rice yield loss (Listianti et al. 2019). However, the yield per hectare from the biobactericide application treatment was not higher than the description of the Ciherang variety, which yields 5-7 tons per hectare. Meanwhile, the production yield in the P1 treatment resulted in a yield of 4.5 tons per hectare. This indicates that disease intensity affects production yield, because bacterial leaf blight attacks the leaves of the plant. Infected leaves cannot produce panicles, resulting in low productivity.

Biological agents have the ability to bind nitrogen, produce plant growth hormones, and dissolve phosphate, among other methods to stimulate growth and increase crop yields. The number of panicles in rice plants is influenced by the nutrient phosphate (Mindari et al. 2018). The panicle is where the grain is attached, and its length determines how many panicles or grains are produced on each panicle. The longer the panicle, the more grains are produced on each panicle.

Weight of dry paddy (GKS) per panicle

The test results showed that the *B. subtilis* bactericide treatment had no significant effect on the weight of dry rice grains per panicle. Table 4 shows that there is no significant effect between the treatment of the biobactericide *B. subtilis* and the treatment of the citronella plant pesticide on the weight of dry rice grains per panicle. The weight of dry rice grains per panicle has a difference of only about 3.54%. The attack of the bacteria Xoo which causes bacterial leaf blight is thought to inhibit the development of rice plants. By reducing the

number of growing panicles or preventing rice grains from filling, the fewer leaves have an indirect effect in limiting crop yields (Khaeruni et al. 2014; Kone et al. 2025). The amount of assimilates spread from the source to the hole is indicated by the overall grain content. According to Pranata and Kurniasih (2019), the absence of a percentage of filled grains indicates that the assimilates distributed during filling are rather low.

Table 4. Weight of dry paddy grain (GKS) per panicle

Treatment	Weight of Dry Paddy Grain (g)
Citronella Botanical Pesticide	2.26a
<i>B. subtilis</i>	2.34a

Remarks: Numbers followed by the same letter in the same column show no significant difference according to the Mann-Whitney test ($P > 0.05$).

Table 4 shows that there is no significant effect between the treatment of the biobactericide *B. subtilis* and the treatment of the citronella plant pesticide on the weight of dry rice grains per panicle. The weight of dry rice grains per panicle has a difference of only about 3.54%. The attack of the bacteria Xoo which causes bacterial leaf blight is thought to inhibit the development of rice plants. By reducing the number of growing panicles or preventing rice grains from filling, the fewer leaves have an indirect effect in limiting crop yields (Khaeruni et al. 2014; Kone et al. 2025). The amount of assimilates spread from the source to the hole is indicated by the overall grain content. According to Pranata and Kurniasih (2019), the absence of a percentage of filled grains indicates that the assimilates distributed during filling are rather low.

Grain weight per 1000 grains

Paddy quality is determined by weighing 1,000 grains; the heavier the grain, the higher the quality and the more substantial the grain. The results of the average dry paddy weight test per 1,000 grains are presented in Table 5.

Table 5. Weight of 1000 grains of dry paddy from rice fields

Treatment	Weight of 1000 Grains (g)
Citronella Botanical Pesticide	29.47a
<i>B. subtilis</i>	32.56b

Remarks: Numbers followed by different letters in the same column indicate a significant difference according to the Paired Samples T Test ($P < 0.05$).

Table 5 shows the results of observations of the *B. subtilis* biobactericide treatment, which significantly affected the weight of dry paddy rice per 1000 grains. The weight of 1000 grains of full paddy in the *B. subtilis* treatment was 32.56 grams; this weight is in accordance with the description of the Ciherang variety. This condition is caused by the use of *B. subtilis* bacteria, which produce growth regulators that have the ability to stimulate root development (Agustiansyah et al. 2020).

B. subtilis treatment had a significant effect on the weight of paddy per 1000 grains. This indicates that the photosynthesis process in plants is functioning well, resulting in the production of assimilates which are the final product (Pranata and Kurniasih 2019). In addition to producing plant hormones, vitamins, and amino acids, *B. subtilis* also functions as N binding and absorbing P and K (Sitepu et al. 2017). Bacteria take nutrients from available nutrients for their own growth and development, thereby slowing plant growth and development and reducing fruit production. Apart from that, the concentration given is low and the application time is too long, making it difficult to synthesize growth regulators (Nusyirwan and Syahadah 2020).

This is suspected to be due to the attack of Xoo, which causes bacterial leaf blight, which will inhibit growth. This research is supported by Pranata and Kurniasih (2019), who stated that reducing the number of leaves definitely reduces production by reducing the number of leaves formed, thus reducing grain filling. The extent of assimilates distribution from the source to the hole is indicated by the overall grain content. The absence of a total percentage of filled grains indicates that the assimilates dispersed during filling are quite low.

CONCLUSIONS AND SUGGESTIONS

The biological agents *B. subtilis* was able to inhibit the spread of bacterial leaf blight disease. The spread of bacterial leaf blight disease attacks will be inhibited because *B. subtilis* bacteria are effective as antagonists of the pathogen Xoo with a medium inhibitory power category with a suspension density of antagonistic bacteria 33×10^4 cfu.ml⁻¹. The treatment of *B. subtilis* biobactericide and citronella plant pesticide showed a significant effect on the intensity of bacterial leaf blight disease namely in the *B. subtilis* treatment 14.72% and citronella plant pesticide 28.97%. The dry weight of rice fields per clump in the *B. subtilis* biobactericide treatment with an average of 27.26 grams was significantly different compared to the citronella plant pesticide treatment with an average of 24.44 g.

ACKNOWLEDGMENTS

The authors would like to express their gratitude to colleagues and other parties involved directly or indirectly in this research. We also extend our gratitude to the Food Crop Production Technology Study Program, Department of Agricultural Production, Politeknik Negeri Jember

REFERENCES

- Agustiansyah A, Ilyas S, Sudarsono S, Machmud M. 2013. Karakterisasi rizobakteri yang berpotensi mengendalikan bakteri *Xanthomonas oryzae* pv. *oryzae* dan meningkatkan pertumbuhan tanaman padi. J Hama dan Penyakit Tumbuh Trop. 13(1):42–51. <https://doi.org/10.23960/j.hptt.11342-51>.
- Agustiansyah A, Ilyas S, Sudarsono S, Machmud M. 2020. Pengaruh perlakuan benih dengan agens hayati dan pemupukan terhadap pertumbuhan tanaman, produksi dan mutu benih padi di lapang. J Agrotropika. 17(2).
- Ali DFI, El-Nahrawy S, EL-Zawawy HAH, Omara AE-D. 2025. Effective applications of *Bacillus subtilis* and *B. amyloliquefaciens* as biocontrol agents of damping-off disease and biostimulation of tomato plants. Stresses. 5(1):9. <https://doi.org/10.3390/stresses5010009>.
- Azman N, Sijam K, Hata E, Othman R, Saud H. 2017. Screening of bacteria as antagonist against *Xanthomonas oryzae* pv. *oryzae*, the causal agent of bacterial leaf blight of paddy and as plant growth promoter. J Exp Agric Int. 16(4):1–15. <https://doi.org/10.9734/JEAI/2017/33697>.
- [BPS Jember] Badan Pusat Statistik Kabupaten Jember. 2023. Kabupaten Jember dalam angka 2023. Jember (ID): Badan Pusat Statistik Kabupaten Jember.
- Barimbing AF. 2022. Uji ekstrak daun sirih (*Piper betle* L.), biji mahoni (*Swietenia mahagoni* (L.) Jacq.) dan daun serai wangi (*Cymbopogon nardus*) dalam mengendalikan *Xanthomonas oryzae* pv. *oryzae* penyebab penyakit hawar daun bakteri pada ta.
- Bota W, Martosupono M, Rondonuwu FS. 2015. Potensi senyawa minyak sereh wangi (citronella oil) dari tumbuhan *Cymbopogon nardus* L. sebagai agen antibakteri BT - Prosiding Semnastek.
- Djarmiko HA, Kurniawan DU, Prihatiningsih N. 2022. Potential of *Bacillus subtilis* potato isolate as biocontrol agent of *Xanthomonas oryzae* pv. *oryzae* and candidate for nanosuspension formula. Biodiversitas J Biol Divers. 23(7):3313–3317. <https://doi.org/10.13057/biodiv/d230701>.
- Donggulo C V, Lapanjang IM, Made U. 2017. Pertumbuhan dan hasil tanaman padi (*Oryza sativa* L.) pada berbagai pola jarak legowo dan jarak tanam. Agrol J Ilmu-Ilmu Pertan. 24(1):27–35.
- Erdiansyah I, Syarief M, Kusairi MI. 2021. The effect of color type and light intensity of Light Emitting Diode (LED) light traps on the types and number of pest insect catches in rice plantations. Cropsaver. 4(1):10–14. <https://doi.org/10.24198/cropsaver.v4i1.28555>.
- [Kementan] Direktorat Jenderal Tanaman Pangan. 2018. Petunjuk teknis pengamatan dan pelaporan organisme pengganggu tumbuhan dan dampak perubahan iklim (OPT-DPI) [revisi 2015]. Jakarta (ID): Kementerian Pertanian RI.
- Khaeruni A, Taufik M, Wijayanto T, Johan E. 2014. Perkembangan penyakit hawar daun bakteri pada tiga varietas padi sawah yang diinokulasi pada beberapa fase pertumbuhan. J Fitopatol Indones. 10(4):119–125. <https://doi.org/10.14692/jfi.10.4.119>.
- Kone S, Konate K, Barro A, Ouedraogo A, Wonni I. 2025. New rice lines resistant to *Xanthomonas oryzae* pv. *oryzae*: a Promising pathway to food security in Burkina Faso. Adv Agric Technol Plant Sci. 8(2):1–14. <https://doi.org/10.63235/AATPS.180176>.
- Listianti NN, Winarno W, Erdiansyah I. 2019. Pemanfaatan ekstrak daun pepaya (*Carica papaya* L.) sebagai insektisida nabati pengendali walang sangit

- (*Leptocorisa acuta*) pada tanaman padi. Agriprima J Appl Agric Sci. 3(1):81–85. <https://doi.org/10.25047/agriprima.v3i1.142>.
- Meimaharani R, Listyorini T. 2013. Analisis varian (ANOVA) untuk mengetahui statistik tingkat kemajuan prestasi karate di Kabupaten Kudus. In: Abidin Z, Subhan S, Triharto WB, editors. *Cloud computing security*. Prosiding seminar nasional ilmu komputer 2013; cloud 23 November 2013, ID Semarang. Semarang (ID): Jurusan Ilmu Komputer, FMIPA UNNES. p. 9–11.
- Mindari W, Widjajani BW, Priyardarsini R. 2018. Kesuburan tanah dan pupuk. Yogyakarta: Gosyen Publishing.
- Molotoks A, Smith P, Dawson TP. 2021. Impacts of land use, population, and climate change on global food security. Food Energy Secur. 10(1). <https://doi.org/10.1002/fes3.261>.
- Moningka CNG, Ludong DPM, Rumambi DP. 2020. Kajian irigasi mikro pada sistem hidroponik padi (*Oryza sativa* L.) varietas serayu dalam rumah tanaman. J Teknol Pertan (Agricultural Technol Journal). 11(1). <https://doi.org/10.35791/jteta.11.1.2020.29981>.
- Nurkartika R, Ilyas SIS, Machmud M. 2017. Aplikasi agens hayati untuk mengendalikan hawar daun bakteri pada produksi benih padi. J Agron Indones (Indonesian J Agron. 45(3):235–242.
- Nusyirwan N, Syahadah RA. 2020. Pengaruh bakteri endofit *Bacillus subtilis* dalam upaya meningkatkan hasil pertumbuhan dan produksi pada tanaman cabai merah (*Capsicum annuum* L.). J Biosains. 6(2):53. <https://doi.org/10.24114/jbio.v6i2.15219>.
- Pandango S, Widnyana IK, Sapanca PLY. 2018. Pengaruh lama perendaman benih dengan bakteri *Bacillus spp.* terhadap pertumbuhan dan hasil tanaman kacang tanah. Agrimeta J Pertan Berbas Keseimbangan Ekosist. 8(16):50–55.
- Pranata M, Kurniasih B. 2019. Pengaruh pemberian pupuk kompos jerami padi terhadap pertumbuhan dan hasil padi (*Oryza sativa* L.) pada kondisi salin (Effect of rice straw compost rate on the growth and yield of rice (*Oryza sativa* L.) under saline conditions). 8(2).
- Serrão CP, Ortega JCG, Rodrigues PC, de Souza CRB. 2024. <I>Bacillus</I> species as tools for biocontrol of plant diseases: A meta-analysis of twenty-two years of research, 2000–2021. World J Microbiol Biotechnol. 40(4):110. <https://doi.org/10.1007/s11274-024-03935-x>.
- Sitepu RB, Anas I, Djuniwati S. 2017. Pemanfaatan jerami sebagai pupuk organik untuk meningkatkan pertumbuhan dan produksi padi (*Oryza sativa*) (Utilization of straw as organic fertilizer to increase growth and production of rice (*Oryza sativa*)). Bul Tanah dan Lahan. 1:100–108.
- Srinivas B, Patil VA, Shinde CU, John P, Garde YA, Waghunde RR. 2024. Effect of antagonists and botanicals against *Xanthomonas oryzae* pv. *oryzae* in vitro. Int J Econ Plants. 11(1):7–11. <https://doi.org/10.23910/2/2024.5088c>.
- Sutarman. 2017. Dasar-dasar ilmu penyakit tanaman. Umsida Press.
- Wakil W, Kavallieratos NG, Eleftheriadou N, Ghazanfar MU, El-Shafie HAF, Blankson A, Gurr GM, Dáttilo W, González-Tokman D, Hernández A, et al. 2025. Climate change consequences for insect pest management, sustainable agriculture and food security. Entomol Gen. 45(1):37–51. <https://doi.org/10.1127/entomologia/2024/2774>.
- Wartono G, Mutaqin KH. 2015. Efektivitas formulasi spora *Bacillus subtilis* B12 sebagai agen pengendali hayati penyakit hawar daun bakteri pada tanaman padi. J Penelit Pertan Tanam Pangan. 34(1):21–28.