



## Optimization of Empty Fruit Bunch Application to Improve Soil Organic Carbon, Total Soil Nitrogen, Earthworm Populations, and Oil Palm Performance on Spodosols

Adhy Ardiyanto\*, Suria Darma Tarigan, Rahayu Widyastuti, Budi Nugroho and Fathan Aldi Rivai

Department of Soil Science and Land Resources, Faculty of Agriculture, IPB University, Bogor, Indonesia

\*Corresponding author: 198ladhy@apps.ipb.ac.id

### Abstract

Using oil palm empty fruit bunches (EFB) as an organic amendment is a crucial strategy to improve soil quality in oil palm plantations, particularly on nutrient-poor Spodosol soils. This study aimed to evaluate the effects of 2 EFB application methods (incorporation into the soil and surface mulching) on soil organic carbon (SOC), nitrogen content, earthworm populations as soil biological indicators of fertility, and oil palm performance. A 12-month field experiment was conducted with observations every 4 months, using a completely randomized design (CRD) with 3 treatments and 9 replications each ( $n = 27$  experimental units). The treatments included no EFB application (T0), EFB incorporated into the soil (T1), and EFB applied as surface mulch (T2). The results showed that both EFB application methods significantly increased SOC compared to the control, with the highest SOC observed in T2 ( $2.30 \pm 0.34\%$ ) after 12 months. Earthworm populations also increased progressively, reaching 19.33 individuals  $m^{-2}$  in T2. Pearson correlation analysis revealed a strong positive relationship between SOC content and earthworm abundance ( $r = 0.60$ ;  $p < 0.001$ ). These results confirm that EFB application, especially through surface mulching, is an effective and sustainable approach to enhancing soil quality, biological activity, and ecosystem functioning on Spodosols in oil palm plantations.

**Keywords:** EFB mulching; organic amendment; *Pontoscolex corethrurus*; soil biological indicators; Spodosols

**Cite this as:** Ardiyanto, A., Tarigan, S. D., Widyastuti, R., Nugroho, B., & Rivai, F. A. (2026). Optimization of Empty Fruit Bunch Application to Improve Soil Organic Carbon, Total Soil Nitrogen, Earthworm Populations, and Oil Palm Performance on Spodosols. *Caraka Tani: Journal of Sustainable Agriculture*, 41(2), 199-215. doi: <http://dx.doi.org/10.20961/carakatani.v41i2.110106>

### INTRODUCTION

Oil palm is one of the leading plantation commodities in Indonesia, with strong competitiveness and a promising future as a sustainable industry. According to data from Statistics Indonesia (2025), the area of oil palm plantations increased by 203.07% between 2003 and 2024. Oil palm plantations contribute to job creation and regional economic growth (Syahza and Asmit, 2019). Over recent decades, oil palm expansion has increasingly extended into marginal lands, including Spodosols that are widely distributed across Borneo Island. Spodosols are characterized by a spodic horizon

formed through podzolization, involving the accumulation of organic matter complexed with aluminum and iron in subsurface layers. They typically occur in coarse-textured, quartz-rich soils with low organic matter content (Huang et al., 2022).

Intensive eluviation–illuviation processes in sandy parent materials create a pronounced contrast between the eluvial and spodic horizons, as reported in several recent studies (Ferro-Vázquez et al., 2020; Souza Junior et al., 2023; Fedenko et al., 2024; Kuligiewicz et al., 2024; Petticord et al., 2025). The characteristics of

\* Received for publication October 19, 2025

Accepted after corrections March 8, 2026

Spodosols promote high nutrient leaching and limited macronutrient availability, resulting in low oil palm productivity without appropriate nutrient management, as inorganic fertilizer efficiency in sandy soils is constrained by rapid leaching losses (Abdelfattah et al., 2021). Therefore, integrated nutrient management approaches combining organic and inorganic inputs are more relevant (Wang et al., 2024).

The rapid growth of the oil palm industry has also generated large amounts of biomass waste, particularly oil palm empty fruit bunches (EFB), which are expected to continue increasing as palm oil production rises (Januari et al., 2020). EFB is the main residue from fresh fruit bunch (FFB) processing and accounts for approximately 23% of the total harvested biomass, with high lignocellulosic and organic matter content (Sari et al., 2018). Despite its slow decomposition, EFB supplies carbon and nutrients for gradual-release soil amendment (Supriatna et al., 2022; Blankson et al., 2025), yet it remains underutilized and is often stockpiled or burned, contributing to greenhouse gas emissions and air pollution (Batcha et al., 2020).

Intensive oil palm management, involving frequent weeding and biomass removal without adequate residue return, can accelerate soil organic matter decomposition, leading to long-term carbon loss (Ashton-Butt et al., 2018; Formaglio et al., 2020; Rahman et al., 2021). Declines in SOC negatively affect soil structure, water retention, and soil biodiversity, ultimately increasing the risk of land degradation (Ashton-Butt et al., 2019; Kaupper et al., 2020). SOC is a key indicator of soil quality and fertility, contributing to aggregate formation, water storage, and energy supply for soil organisms (Guo et al., 2025; Santos et al., 2025).

Higher SOC content supports the development of more stable microhabitats for soil fauna, including earthworms, which are recognized as ecosystem engineers (Ahmed and Al-Mutairi, 2022). Earthworms enhance soil porosity and nutrient cycling through bioturbation activities and the production of nutrient-rich casts (Hallam et al., 2020; Capowiez et al., 2021; Traoré et al., 2022; Amadou et al., 2025). The addition of organic materials has been shown to increase earthworm populations by providing food resources and improving soil habitat conditions (Panta et al., 2026).

The use of EFB has been widely studied for various purposes, including bioenergy production and industrial materials (Mahardika et al., 2024;

Utomo et al., 2024; Chitraningrum et al., 2025; Rosa Silva et al., 2026). In situ EFB application in oil palm plantations offers strategic advantages by reducing transport costs and carbon emissions while returning organic matter to the system. Several studies have examined the relationship between EFB application and earthworm populations, but most of these investigations were conducted outside the context of Spodosols, leaving a limited understanding of their interactions in sandy marginal soils (Tarigan et al., 2019; Hau et al., 2020; Lew et al., 2020; Edyson et al., 2022).

Organic matter application is a key soil management practice that increases SOC, improves soil physical and chemical properties, and supports biological activity, all of which are critical for agricultural productivity (Chiodi et al., 2025; Xu et al., 2025). The effectiveness of organic amendments depends on the application method, as soil incorporation and surface mulching produce distinct decomposition, carbon, and nitrogen dynamics (Readyhough et al., 2021; Abbas et al., 2024; Maticic et al., 2024). Soil health is fundamentally influenced by biological components, as biological activity drives improvements in soil physical and chemical properties (Zhelezova et al., 2025). Soil fauna play an important role in organic matter decomposition and nutrient cycling, making them reliable indicators of soil quality (Zhang et al., 2023; Mamabolo et al., 2024).

Earthworm populations serve as biological indicators of soil responses to EFB application, enhancing decomposition and nitrogen availability, which are essential for oil palm growth (Yadegari et al., 2020; Arifin et al., 2022; De la Peña et al., 2023). Nitrogen derived from EFB, which has a relatively high C/N ratio, does not necessarily increase soil nitrogen levels because some may be immobilized by microbes or be rapidly taken up by plants (Hu and Zhu, 2025; van Rijssel et al., 2025). Therefore, evaluating the effectiveness of EFB application requires consideration of both soil and plant nitrogen status, as represented by leaf nitrogen content.

Existing studies on oil palm EFB management have generally focused on short-term nutrient dynamics and yield responses, with limited attention to comparative effects of application methods on biological soil indicators and long-term SOC dynamics, particularly in Spodosols. Interactions among EFB application methods, SOC, earthworm populations, and soil and plant nitrogen status under field conditions remain

insufficiently documented. Accordingly, the novelty of this study lies in its field-based comparative approach that integrates chemical, biological, and plant-based indicators to evaluate the effectiveness of different EFB application methods on Spodosols. Based on these considerations, this study aims to analyze changes in SOC following EFB application through soil incorporation and surface mulching, evaluate earthworm population responses, and assess their effects on soil nitrogen, leaf nitrogen content, and oil palm productivity.

## MATERIALS AND METHOD

### Research location and time

The study was conducted in a PT. Bumitama Gunajaya Agro (BGA) oil palm plantation on Spodosol soil in Kotawaringin Timur Regency (112°07'29" to 113°14'22" E and 1°11'50" to 3°18'51" S), Central Kalimantan, Indonesia (Borneo Island). The field experiment lasted for 12 months, from August 2023 to August 2024. Soil sampling and earthworm population observations were conducted every 4 months to monitor temporal changes. Soil sample analyses were conducted at the PT. BGA analytical laboratory.

### Materials and equipment

The main material used in this study was EFB obtained from palm oil mill waste near the research site. Spodosol soil was used as the natural growing medium, while oil palm plants (*Elaeis guineensis* Jacq.) in the production phase (planted in 2011) served as the research objects. The tools used included a mini excavator (PC 50),

soil sample rings, a soil auger, a sharp field knife, a hoe, and a measuring ruler. Soil fauna were identified using hand-sorting monoliths (50 cm × 50 cm × 20 cm), and organic carbon was analyzed using the Walkley and Black method.

### Experimental design

The experiment was arranged in a completely randomized design (CRD) with 3 treatments and 9 replications, yielding 27 experimental plots. Each plot consisted of approximately 25 oil palm trees, which were treated as subsampling units. The treatments included: T0 = a control without EFB application; T1 = the application of 200 kg EFB per tree incorporated into the soil on both sides of the plant within a 2 m radius from the trunk; and T2 = the application of 200 kg EFB per tree applied as surface mulch within a 2 m radius from the trunk. Soil samples were collected at 2 depths (0 to 20 cm and 20 to 40 cm) and at 2 sampling locations: point X, corresponding to the EFB application area, and point Y, located 50 cm from point X toward the oil palm trunk. The different locations of points X and Y are illustrated in Figure 1.

### Soil sampling and SOC analysis

SOC property sampling was conducted in August 2023 (before treatment) and every 4 months (December 2023, April 2024, and August 2024). Soil samples were collected from a depth of 0 to 20 cm at points X and Y to determine SOC, then composited and homogenized for analysis of earthworm populations and soil nitrogen. SOC content was determined using the Walkley and Black method (Nelson and Sommers, 1982). This method involves oxidizing organic matter




Figure 1. Location of soil sampling points

with potassium dichromate ( $K_2Cr_2O_7$ ) under acidic conditions, followed by titration with ferrous sulfate ( $FeSO_4$ ). The results were used to evaluate the impact of EFB application on SOC content. Total soil nitrogen was determined using the Kjeldahl method, in which finely ground soil samples were digested with concentrated  $H_2SO_4$  and a catalyst mixture, followed by distillation and titration to quantify total nitrogen content.

### Earthworm population calculation

Earthworm populations were observed concurrently with soil sampling in each experimental unit. Earthworms were collected using the hand-sorting method on a soil monolith measuring 50 cm × 50 cm × 20 cm, which represented the EFB application zone. All earthworm individuals found were manually separated from the soil, then counted and expressed as the number of individuals per square meter (individuals  $m^{-2}$ ) as a biological indicator of soil quality. The use of the hand-sorting method followed the recommendation of Nadolny et al. (2020), which is commonly used in soil ecology studies to obtain accurate field estimates of earthworm populations.

### Leaf sampling

Leaf nitrogen concentration was determined using the Kjeldahl method following the procedure described by Kamireddy et al. (2023). Fully expanded leaf samples (fronds 17) from 13-year-old oil palm trees were collected in August 2023 and subsequently at 4-month intervals for up to 12 months. The samples were oven-dried at 65 °C to constant weight, finely ground, digested with concentrated  $H_2SO_4$ , and distilled to determine total nitrogen content.

### Measurement of oil palm productivity

Oil palm productivity was assessed based on FFB yield data collected over a 6-month harvest period (September 2024 to February 2025), commencing 1 year after treatment application. FFB were harvested at 7 to 8-day intervals from 27 observation plots (9 palms per plot), and the yield data were subsequently converted to tons per hectare per 6-month period.

### Data analysis

Data on SOC content, earthworm populations, soil nitrogen content, and leaf nitrogen content were analyzed using analysis of variance (ANOVA) according to the applied experimental design. ANOVA is a statistical method used to test whether there are significant differences among treatment means for the observed variables

(Chicco et al., 2025). When a significant effect was detected at the 5% confidence level, Tukey's honestly significant difference (HSD) post hoc test was applied to determine which treatment pairs differed significantly. The correlation between SOC content and earthworm populations was analyzed using Pearson's correlation. Pearson's correlation is a parametric statistical analysis used to measure the strength and direction of a linear relationship between 2 quantitative variables (Tianle et al., 2026).

## RESULTS AND DISCUSSION

### Effect of EFB application on SOC

SOC content increased clearly in the EFB treatments, especially under surface mulching, while SOC in the control treatment tended to remain stable or decrease (Figure 2). At the 0 to 20 cm soil depth, SOC increased by 0.54% with surface application and by 0.48% with incorporated application, showing the effectiveness of EFB as a carbon source. Incorporation of EFB into the soil also increased SOC, but the increase was slightly lower due to more humid conditions (Tao et al., 2018). At depths of 20 to 40 cm, the increase was smaller because dissolved organic matter movement was limited by the characteristics of Spodosols. However, the moderate increase in the EFB treatments still indicates the contribution of dissolved organic matter to subsoil carbon. Noirod et al. (2022) reported that applying EFB in oil palm plantations increases SOC content compared to no treatment, mainly due to its role as a mulch in maintaining soil moisture and temperature, which supports decomposer activity. Rosenani et al. (2016) also stated that EFB decomposition occurs gradually and contributes to the addition of organic matter within 12 to 18 months after application.

The faster increase of SOC in T2 indicates the contribution of more easily decomposed organic matter fractions, while T1 shows slower but more stable carbon accumulation. This is consistent with Basile-Doelsch et al. (2020), who explained that organic carbon from fresh residues undergoes gradual humification and forms more protected organo-mineral associations in deeper layers. The incorporation of EFB by burying can increase SOC content (Nyasapoh et al., 2026); however, other studies have confirmed that surface application still produces the highest SOC due to better retention of biological activity (Adu et al., 2022).




Figure 2. Dynamics of SOC under EFB treatments at different soil depths and observation locations: a) Point X (0–20 cm); b) Point X (20–40 cm); c) Point Y (0–20 cm); and d) Point Y (20–40 cm)

In terms of location, SOC values from August 2023 to August 2024 showed that site X (the EFB application spot) consistently had higher SOC than site Y, located 50 cm from the application point. This indicates that the improvement in soil organic matter from EFB is highly localized around the application zone. EFB has coarse particles and decomposes slowly, so lateral carbon movement is limited within 1 year (Shumba et al., 2024). These findings align with Rosenani et al. (2016), who reported that carbon accumulation remains concentrated in the organic residue application point. Therefore, the EFB application strategy should consider spacing and distribution patterns to ensure soil organic matter improvement across a wider plantation area.

SOC at 0 to 20 cm was higher than that at 20 to 40 cm for all treatments. This pattern reflects the concentration of organic inputs in the surface soil, where residue decomposition and microbial processes are generally more intense (Brindis-Santos et al., 2021; Wang et al., 2025). However, SOC increase at 20 to 40 cm was still observed at site X, indicating the contribution of dissolved organic carbon moving downward through water percolation and bioturbation (Rizinjirabake et al., 2019). The increase in deeper layers occurred more slowly because organic matter movement takes more time and is influenced by carbon stabilization processes through interactions with soil minerals such as Al and Fe oxides (Basile-Doelsch et al., 2020; Burgeon et al., 2021). These

mechanisms are associated with the formation of organo-mineral complexes, which contribute to long-term carbon storage but only when there is a continuous supply of organic matter (Shabtai et al., 2024). Thus, repeated and long-term EFB application has the potential to increase carbon stocks not only in the topsoil but also in deeper layers through gradual humification, vertical carbon transport, and mineral stabilization.

Although incorporation of EFB into the soil (T1) increases contact between organic matter and the soil matrix and stimulates microbial activity for faster decomposition, this study showed that surface application (T2) tended to produce slightly higher SOC in the upper soil layer. This effect is related to mulching, which maintains soil moisture, reduces temperature fluctuations, and slows carbon mineralization, resulting in more stable carbon accumulation (Khoramizadeh et al., 2021; Rudolf et al., 2021). These findings show that surface mulching is more effective for long-term SOC enhancement than incorporation.

The decrease in SOC observed in treatment T1 at 20 to 40 cm depth in December 2023 ( $0.88 \pm 0.39\%$ ) was influenced by the positive soil priming effect, where the addition of fresh organic matter stimulates microbial activity and accelerates the decomposition of older, more resistant soil carbon (Mazzilli et al., 2014; Bernard et al., 2022). This mechanism may temporarily reduce carbon stock due to oxidation of older carbon to  $\text{CO}_2$ , before the balance is restored through humification and stabilization of newly added carbon associated with soil minerals (Fang et al., 2023; Martial et al., 2023). Therefore, the EFB application may temporarily accelerate the decomposition of old carbon before stabilization occurs.

Temporally, SOC dynamics showed the decomposition phases of EFB biomass. The increase from August to December 2023 reflected the rapid use of easily decomposed fractions as microbial energy sources. From December 2023 to April 2024, SOC tended to stagnate or decrease because the easily decomposed fractions were depleted, shifting decomposition to more resistant

lignocellulose components. The minor fluctuation from April to August 2024 reflects the gradual breakdown of lignin and cellulose as the dominant components of EFB, which slows the decomposition rate but contributes to long-term stable carbon formation.

### Effect of EFB application on total nitrogen in soil and oil palm leaves

Total soil nitrogen is one of the important indicators of soil fertility because it reflects the long-term nitrogen reserves stored in soil organic matter. In Spodosols, which generally have relatively low nutrient content and are dominated by stable organic matter, nitrogen dynamics are strongly influenced by organic matter inputs. The results of this study show that total soil nitrogen content was affected by oil palm EFB treatments and observation time, indicating the occurrence of nitrogen transformation processes in the soil (Table 1). At the initial observation (August 2023), soil nitrogen content across all treatments was relatively uniform, with values ranging from 0.09 to 0.11. The treatment without EFB (T0) recorded a value of  $0.11 \pm 0.02$ , while T1 and T2 showed values of  $0.09 \pm 0.00$  and  $0.10 \pm 0.02$ , respectively. These results indicate that before the treatments produced a significant effect, soil nitrogen status among treatments was still at a relatively similar level.

At the December 2023 observation, soil nitrogen content decreased in all treatments, with a more pronounced decline in treatments T1 and T2, which showed values of  $0.08 \pm 0.01$  and  $0.08 \pm 0.02$ , respectively. In contrast, treatment T0 maintained a slightly higher value of  $0.09 \pm 0.02$ . The decrease in soil nitrogen in treatments receiving EFB indicates the occurrence of an early phase of nitrogen immobilization. Nitrogen immobilization is a condition in which mineral nitrogen in the soil is used by microorganisms to support the decomposition of organic materials with a relatively high C/N ratio (Cao et al., 2021). At this stage, nitrogen is not lost from the system but is temporarily bound in microbial biomass, resulting in lower measured soil nitrogen levels.

Table 1. Soil nitrogen content across different treatments and observation times

Treatment	Soil nitrogen content			
	August 2023	December 2023	April 2024	August 2024
T0	$0.11 \pm 0.02^{bcd}$	$0.09 \pm 0.02^{cd}$	$0.11 \pm 0.03^{bcd}$	$0.17 \pm 0.05^a$
T1	$0.09 \pm 0.00^{cd}$	$0.08 \pm 0.01^d$	$0.07 \pm 0.03^d$	$0.15 \pm 0.03^{ab}$
T2	$0.10 \pm 0.02^{cd}$	$0.08 \pm 0.02^d$	$0.10 \pm 0.03^{bcd}$	$0.13 \pm 0.04^{abc}$

Note: Values are means  $\pm$  SE. Different letters in the same row indicate significant differences (Tukey's test,  $\alpha = 0.05$ )

Differences among treatments became more evident at the April 2024 observation. Treatment T1 had the lowest soil nitrogen content ( $0.07\pm 0.03$ ), which was significantly lower than the other treatments. Meanwhile, treatments T0 and T2 recorded values of  $0.11\pm 0.03$  and  $0.10\pm 0.03$ , respectively. The low soil nitrogen content in treatment T1 indicates that nitrogen released from EFB decomposition had not yet accumulated as measurable soil nitrogen but remained bound within the organic fraction (Wierzbowska et al., 2021) and had undergone turnover through soil microbial activity (Bossolani et al., 2023). Because burying EFB increases contact between organic material and soil, it enhances microbial activity and prolongs the nitrogen immobilization phase.

At the final observation (August 2024), soil nitrogen content increased in all treatments. Treatment T0 showed the highest value ( $0.17\pm 0.05$ ), and was significantly higher than the other treatments. Treatments T1 and T2 reached  $0.15\pm 0.03$  and  $0.13\pm 0.04$ , respectively. This increase indicates that nitrogen mineralization had occurred. However, soil nitrogen content in the EFB treatments remained lower than in the control, suggesting that nitrogen released through mineralization is generally rapidly taken up by plants and soil microorganisms and therefore does not always accumulate as total soil nitrogen (Ward and Bradford, 2025).

Changes in oil palm leaf nitrogen content indicate plant responses to soil nitrogen dynamics resulting from EFB application. Leaf nitrogen content reflects the nitrogen that is absorbed and used by the plant and therefore does not always correspond directly with total soil nitrogen content. The results of the leaf nitrogen observations are presented in Table 2. At the initial observation (August 2023), leaf nitrogen content was relatively uniform across all treatments, with values ranging from 2.15 to 2.24%. Treatment T0 recorded a value of  $2.15\pm 0.06\%$ , while treatments T1 and T2 showed values of  $2.20\pm 0.08\%$  and  $2.24\pm 0.08\%$ ,

respectively. These values reflect the initial nitrogen status of the plants before the EFB treatment had an effect.

At the December 2023 observation, leaf nitrogen content increased in all treatments, with the highest value recorded in treatment T2 ( $2.51\pm 0.22\%$ ), followed by T1 ( $2.45\pm 0.05\%$ ) and T0 ( $2.42\pm 0.18\%$ ). This increase indicates that part of the nitrogen released during decomposition and mineralization had become available and was absorbed by the plants. Although soil nitrogen content in the EFB treatments was relatively lower during this period, available nitrogen appeared to be utilized more rapidly by plants rather than accumulated in the soil. The most pronounced increase in leaf nitrogen content occurred at the April 2024 observation, particularly in treatment T1, which reached the highest value of  $2.62\pm 0.10\%$  and was significantly different from the other treatments. The treatment T2 also showed a relatively high value ( $2.50\pm 0.10\%$ ), whereas T0 remained lower ( $2.44\pm 0.17\%$ ). These results indicate that burying EFB improved plant nitrogen uptake efficiency. This improvement may occur through increased nitrogen availability in the root zone (Phillips et al., 2022).

At the final observation (August 2024), leaf nitrogen content decreased in all treatments, with values of  $2.38\pm 0.16\%$  (T0),  $2.33\pm 0.13\%$  (T1), and  $2.28\pm 0.12\%$  (T2), respectively. This decline indicates that leaf nitrogen status is influenced not only by soil nitrogen availability but also by plant physiological factors, such as nitrogen redistribution to other plant organs and changes in nitrogen demand during growth (Li et al., 2025). Pearson correlation analysis in the final study period (August 2024) yielded a significant positive relationship between total soil nitrogen and leaf nitrogen content ( $r = 0.73$ ;  $p = 0.026$ ). These results indicate that the increase in soil nitrogen due to EFB application directly contributed to the improvement in nitrogen status of oil palm plants.

Table 2. Nitrogen content of oil palm leaves (Mean $\pm$ SE) across different treatments and observation times

Treatment	Nitrogen content of oil palm leaves			
	August 2023	December 2023	April 2024	August 2024
T0	$2.15\pm 0.06^d$	$2.42\pm 0.18^{bc}$	$2.44\pm 0.17^b$	$2.38\pm 0.16^{bc}$
T1	$2.20\pm 0.08^{cd}$	$2.45\pm 0.05^b$	$2.62\pm 0.10^a$	$2.33\pm 0.13^c$
T2	$2.24\pm 0.08^c$	$2.51\pm 0.22^{ab}$	$2.50\pm 0.10^{ab}$	$2.28\pm 0.12^c$

Note: Values are means $\pm$ SE. Different letters in the same row indicate significant differences (Tukey's test,  $\alpha = 0.05$ )

### Effect of EFB application on earthworm population

The application of oil palm EFB significantly affected the dynamics of earthworm populations throughout the study period (Table 3). The earthworms identified in this study were *Pontoscolex corethrurus*, belonging to the family Glossoscolecidae. This species possesses an elongated cylindrical body with distinct segmentation and bilateral symmetry (James et al., 2019). *P. corethrurus* is recognized as a tolerant species capable of thriving under a wide range of environmental conditions and can become abundant in habitats with relatively low litter availability (Potapov et al., 2021). It has also been frequently reported in tropical oil palm plantation ecosystems (Azhar et al., 2024; Abdi et al., 2025; Al-Saedi et al., 2025) and tends to dominate earthworm communities compared with other species (Taheri et al., 2018).

In general, the population increased over time in all treatments, with the highest increase consistently observed in treatment T2 (surface-applied EFB). Analysis of variance showed that both treatment and observation time had significant effects on the earthworm population ( $p < 0.05$ ). At the beginning of the observation, populations were relatively low, and there were no significant differences among treatments, reflecting the poor organic matter content of the Spodosols. Significant differences began to emerge in December 2023 and became more apparent in subsequent observations, with the peak population in T2 reaching 19.33 individuals  $m^{-2}$  in August 2024, markedly higher than in T1 and T0. This increase highlights the importance of surface-availability soil organic matter as the primary food source and habitat supporting earthworm activity (Vršič et al., 2021).

The earthworm population showed different responses to the organic material management treatments throughout the observation period. During the first two sampling times (August and December 2023), earthworm populations were still low and not significantly different among treatments. This indicates that soil biota requires

an adaptation period following EFB application before population increases can occur. Significant differences began to appear in April 2024, when treatment T2 caused a clear increase in population compared with T0 and T1. This pattern continued until August 2024, with the highest population observed in T2 ( $19.33 \pm 16.49$  individuals  $m^{-2}$ ). This result shows that placing organic material on the soil surface provides a more favorable habitat, food source, and suitable microhabitat conditions for earthworms.

As a surface mulch, EFB also helps maintain soil moisture, stabilize temperature, and create suitable microhabitats for earthworms (Supriatna et al., 2022). In contrast, in T1, earthworm population increases were slower and not significantly different from the control until the end of the observation. Semi-anaerobic conditions in the subsurface layer are less favorable for earthworm activity and reproduction (Tao et al., 2018). Therefore, T1 did not show significant differences. Meanwhile, in T2, a significant difference in earthworm population was already observed since April 2024.

The earthworm population increased progressively during the observation period, with a significant rise after 8 to 12 months of EFB application, especially in T2. This increase has important implications for soil ecosystem functions. As ecosystem engineers, earthworms improve soil structure through bioturbation, increase water infiltration, accelerate nutrient cycling, and enrich soil organic matter (Ahmed and Al-Mutairi, 2022; Ratsiatosika et al., 2024). In addition, positive interactions with decomposer microorganisms can enhance mineralization efficiency and create ecological feedback that strengthens long-term soil fertility (Gutierrez Al-Khudhairi et al., 2023).

Organic matter decomposition by earthworms leads to the formation of casts, while their digestive activity, soil particle aggregation, and movement redistribute SOC throughout the soil (Iordache, 2023). The process of residue fragmentation and the formation of nutrient-rich casts (worm droppings) encourages nitrogen

Table 3. Earthworm population (individuals  $m^{-2}$ ) across different treatments and observation times

Time	T0 (Control)	T1 (Incorporated)	T2 (Surface)
August 2023	0.74 $\pm$ 1.51 <sup>b</sup>	0.67 $\pm$ 2.00 <sup>b</sup>	0.89 $\pm$ 2.03 <sup>b</sup>
December 2023	1.78 $\pm$ 2.54 <sup>b</sup>	0.67 $\pm$ 1.00 <sup>b</sup>	3.78 $\pm$ 2.91 <sup>b</sup>
April 2024	0.89 $\pm$ 1.45 <sup>b</sup>	3.11 $\pm$ 4.01 <sup>b</sup>	14.44 $\pm$ 6.15 <sup>a</sup>
August 2024	4.44 $\pm$ 2.60 <sup>b</sup>	5.56 $\pm$ 4.22 <sup>b</sup>	19.33 $\pm$ 16.49 <sup>a</sup>

Note: Values are means $\pm$ SE. Different letters in the same row indicate significant differences (Tukey's test,  $\alpha = 0.05$ )

mineralization, thereby increasing the concentration of plant-available nitrogen (Tecimen et al., 2021). These findings confirm that surface spreading of EFB is the most effective strategy to increase earthworm populations and improve organic residue management in oil palm plantation Spodosol soils.

In general, organic carbon content was positively and significantly correlated with earthworm population under EFB treatments, while a weak negative correlation was observed in the control. This confirms that organic material addition plays an important role in enhancing biological activity through positive feedback between organic carbon content and soil biota activity. Increasing SOC supports the growth of microorganisms that serve as food for earthworms, while earthworm activity accelerates organic matter decomposition and nutrient cycling (Guo et al., 2019; Ma et al., 2022).

In T0 ( $r = -0.35$ ;  $p = 0.0337$ ), the weak negative correlation indicates that naturally increasing soil carbon without regulated EFB addition is not sufficient to support earthworm populations. Acidic conditions, low organic matter content, and limited biological activity in Spodosols keep populations low even when carbon content increases slightly. In T1 ( $r = 0.45$ ;  $p = 0.0061$ ), the correlation was positive but weak, suggesting that EFB incorporation has a positive effect on earthworm populations.

In T2 ( $r = 0.60$ ;  $p = 0.0001$ ;  $n = 36$ ), the strong positive correlation indicates a close relationship between the increase in SOC and earthworm population (Figure 3). Surface application of organic material provides direct access for earthworms and helps maintain soil moisture and temperature, thereby creating optimal microhabitat conditions (Radics et al., 2022;

Yaseen and Abdulqadir, 2024). This makes the surface-spreading method the most effective for stimulating earthworm activity and population growth. The differences in correlation strength among treatments show that the effectiveness of increasing earthworm populations is determined not only by the amount of organic material, but also by the application method. The strongest correlation in T2 confirms that applying organic material on the soil surface is the most efficient strategy to improve soil biological functions in Spodosols.

### Effect of EFB application on oil palm productivity

The results of Tukey's HSD post hoc test at the 5% significance level (Table 4) indicate that the application of oil palm EFB had a significant effect on FFB productivity. The treatment without EFB application (T0) resulted in the lowest productivity, at 10.63 tons  $\text{ha}^{-1}$ , and was significantly different from T2. In contrast, the treatment in which EFB was incorporated into the soil (T1) produced an intermediate productivity value (12.80 tons  $\text{ha}^{-1}$ ) and was not statistically different from either T0 or T2, as indicated by the same significance letter (ab). The highest productivity was observed under the T2 treatment, yielding 13.31 tons  $\text{ha}^{-1}$ . The significant difference between T2 and T0 suggests that the presence of EFB generally enhances oil palm productivity; however, its effectiveness is strongly influenced by the application method. These findings emphasize that not only the amount of organic material applied, but also its placement in the field, plays a crucial role in determining crop yield responses.

Treatment T2 is presumed to create more favorable soil conditions for oil palm growth.




Figure 3. Relationship between SOC content and earthworm at a depth of 0-20 cm for T2

Table 4. Post hoc test results for the effect of EFB application on FFB productivity

Treatment	FFB yield (tons ha <sup>-1</sup> )	SE	Significant letter
T0	10.63	±0.63	b
T1	12.80	±0.71	ab
T2	13.31	±0.51	a

Note: Values are means of 6 months of harvest (n = 9). SE = Standard error. Different letters indicate significant differences according to Tukey's HSD test ( $\alpha = 0.05$ )

Organic mulching has been widely reported to reduce evaporation, maintain soil moisture, and improve the physical properties of the topsoil layer (Lopes et al., 2025). These effects are particularly important during dry periods by alleviating water stress and supporting plant physiological processes (Demo and Asefa Bogale, 2024; Vanella et al., 2025). Decomposition of EFB applied on the soil surface occurs relatively more slowly than that of EFB incorporated into the soil. This slower mineralization process allows nutrients to be released gradually and more in line with plant demand. Such synchronization between nutrient availability and plant uptake is believed to contribute to increased FFB productivity (Noirot et al., 2022). Thus, EFB functions not only as a nutrient source but also as a regulator of nutrient dynamics in the soil.

In the T1, decomposition tends to proceed more rapidly due to increased soil microbial activity. This condition may lead to temporary nitrogen immobilization, whereby available soil nitrogen is utilized by microorganisms for organic matter decomposition, thereby reducing nitrogen availability to plants during the early growth stages (Rosenani et al., 2016). This phenomenon may explain why productivity gains under T1 were lower than under T2, although absolute yields remained higher than under T0. From an operational and economic perspective, T2 also offers additional advantages. Surface application of EFB is relatively simpler and requires lower costs compared to incorporation using heavy equipment, such as a PC 50 mini excavator for soil excavation and backfilling as performed in the T1 treatment. By combining the highest productivity and more efficient operational costs, the T2 method provides superior agronomic and economic benefits.

## CONCLUSIONS

The application of EFB improved soil organic carbon levels, increased earthworm abundance, and altered nitrogen dynamics in Spodosols under oil palm cultivation. Surface application of EFB was consistently more effective than

incorporation in stimulating biological activity, promoting carbon stabilization, and increasing nutrient availability. By contrast, incorporation mainly accelerated short-term mineralization processes. Although total soil nitrogen declined temporarily due to microbial immobilization, nitrogen concentrations in oil palm leaves increased, indicating more efficient plant nitrogen uptake. These improvements in soil conditions were reflected in higher FFB yields, particularly under surface-applied EFB. Overall, surface application of EFB can be recommended as a sustainable management strategy to improve soil quality, nutrient cycling, and oil palm productivity, with clear benefits for long-term soil fertility in Spodosols.

## ACKNOWLEDGEMENT

The authors would like to express their highest appreciation to the Management of PT. Bumitama Gunajaya Agro for providing full financial support for this research. Their commitment and contribution were instrumental in enabling the successful completion of this study.

## REFERENCES

- Abbas, A., Naveed, M., Shehzad Khan, K., Ashraf, M., Siddiqui, M. H., Abbas, N., ... & Ali, L. (2024). The efficacy of organic amendments on maize productivity, soil properties and active fractions of soil carbon in organic-matter deficient soil. *Spanish Journal of Soil Science*, 14, 12814. <https://doi.org/10.3389/sjss.2024.12814>
- Abdelfattah, M. A., Rady, M. M., Belal, H. E. E., Belal, E. E., Al-Qthanin, R., Al-Yasi, H. M., & Ali, E. F. (2021). Revitalizing fertility of nutrient-deficient virgin sandy soil using leguminous biocompost boosts *Phaseolus vulgaris* performance. *Plants*, 10(8), 1637. <https://doi.org/10.3390/plants10081637>
- Abdi, A., Sabrina, T., & Sembiring, M. (2025). Earthworm population and diversity in smallholder oil palm plantations under conventional and semi-organic systems in

- Labuhan Batu Regency. *BIO Web of Conferences*, 189, 01015. <https://doi.org/10.1051/bioconf/202518901015>
- Adu, M. O., Atia, K., Arthur, E., Asare, P. A., Obour, P. B., Danso, E. O., ... & Andersen, M. N. (2022). The use of oil palm empty fruit bunches as a soil amendment to improve growth and yield of crops. A meta-analysis. *Agronomy for Sustainable Development*, 42(2), 13. <https://doi.org/10.1007/s13593-022-00753-z>
- Ahmed, N., & Al-Mutairi, K. A. (2022). Earthworms effect on microbial population and soil fertility as well as their interaction with agriculture practices. *Sustainability*, 14(13), 7803. <https://doi.org/10.3390/su14137803>
- Al-Saedi, T. K., Sayed, K., Syakir, M. I., Umar, Y., Azhar, B., Tohiran, K. A., ... & Rahmat, S. R. (2025). Enhancing soil quality and economic value in oil palm plantations: Effects of livestock integration on earthworm abundance and ecosystem services. *Journal for Nature Conservation*, 86, 126939. <https://doi.org/10.1016/j.jnc.2025.126939>
- Amadou, I., Mazurier, A., Caner, L., Zi, Y., Rumpel, C., & Bottinelli, N. (2025). Interactions between earthworm species and soil type influence the porosity of earthworm casts. *Soil Biology and Biochemistry*, 203, 109739. <https://doi.org/10.1016/j.soilbio.2025.109739>
- Arifin, I., Hanafi, M. M., Roslan, I., Ubaydah, M. U., Karim, Y. A., Tui, L. C., & Hamzah, S. (2022). Responses of irrigated oil palm to nitrogen, phosphorus and potassium fertilizers on clayey soil. *Agricultural Water Management*, 274, 107922. <https://doi.org/10.1016/j.agwat.2022.107922>
- Ashton-Butt, A., Aryawan, A. A., Hood, A. S., Naim, M., Purnomo, D., Suhardi, ... & Snaddon, J. L. (2018). Understorey vegetation in oil palm plantations benefits soil biodiversity and decomposition rates. *Frontiers in Forests and Global Change*, 1, 10. <https://doi.org/10.3389/ffgc.2018.00010>
- Ashton-Butt, A., Willcock, S., Purnomo, D., Suhardi, Aryawan, A. A., Wahyuningsih, R., ... & Snaddon, J. L. (2019). Replanting of first-cycle oil palm results in a second wave of biodiversity loss. *Ecology and Evolution*, 9(11), 6433–6443. <https://doi.org/10.1002/ece3.5218>
- Azhar, B., van der Meer, P., Sterenberg, R. F., Yahya, M. S., Razi, N., Burhanuddin, M., ... & Oon, A. (2024). Resilience underground: Understanding earthworm biomass responses to land use changes in the tropics. *Biological Conservation*, 299, 110800. <https://doi.org/10.1016/j.biocon.2024.110800>
- Basile-Doelsch, I., Balesdent, J., & Pellerin, S. (2020). Reviews and syntheses: The mechanisms underlying carbon storage in soil. *Biogeosciences*, 17(21), 5223–5242. <https://doi.org/10.5194/bg-17-5223-2020>
- Batcha, M. F. M., Othaman, M. F., Sabudin, S., Mohammed, A. N., Abdullah, M. K., & Ali, M. F. M. (2020). Combustion and emission of pelletized empty fruit bunch and oil palm shell in a swirling fluidized bed combustor. *Biomass Conversion and Biorefinery*, 10(3), 755–763. <https://doi.org/10.1007/s13399-020-00903-x>
- Bernard, L., Basile-Doelsch, I., Derrien, D., Fanin, N., Fontaine, S., Guenet, B., ... & Maron, P. (2022). Advancing the mechanistic understanding of the priming effect on soil organic matter mineralisation. *Functional Ecology*, 36(6), 1355–1377. <https://doi.org/10.1111/1365-2435.14038>
- Blankson, D., Arthur, E., Atiah, K., Frimpong, K. A., Manfo, P., & Ravnskov, S. (2025). Empty oil palm fruit bunch biochar and compost influence soil properties that drive okra (*Abelmoschus esculentus* L.) nutrient use efficiency and yield. *Soil Science Society of America Journal*, 89(6), e70172. <https://doi.org/10.1002/saj2.70172>
- Bossolani, J. W., Leite, M. F. A., Momesso, L., ten Berge, H., Bloem, J., & Kuramae, E. E. (2023). Nitrogen input on organic amendments alters the pattern of soil–microbe–plant co-dependence. *Science of The Total Environment*, 890, 164347. <https://doi.org/10.1016/j.scitotenv.2023.164347>
- Brindis-Santos, A. I., Palma-López, D. J., Mata-Zayas, E. E., & Palma-Cancino, D. J. (2021). Impacts of oil palm cultivation on soil organic carbon stocks in Mexico: Evidence from plantations in Tabasco State. *Cahiers Agricultures*, 30, 47. <https://doi.org/10.1051/cagri/2021033>
- Burgeon, V., Fouché, J., Leifeld, J., Chenu, C., & Cornéilis, J.-T. (2021). Organo-mineral

- associations largely contribute to the stabilization of century-old pyrogenic organic matter in cropland soils. *Geoderma*, 388, 114841. <https://doi.org/10.1016/j.geoderma.2020.114841>
- Cao, Y., He, Z., Zhu, T., & Zhao, F. (2021). Organic-C quality as a key driver of microbial nitrogen immobilization in soil: A meta-analysis. *Geoderma*, 383, 114784. <https://doi.org/10.1016/j.geoderma.2020.114784>
- Capowiez, Y., Sammartino, S., Keller, T., & Bottinelli, N. (2021). Decreased burrowing activity of endogeic earthworms and effects on water infiltration in response to an increase in soil bulk density. *Pedobiologia*, 85–86, 150728. <https://doi.org/10.1016/j.pedobi.2021.150728>
- Chicco, D., Sichenze, A., & Jurman, G. (2025). A simple guide to the use of Student's t-test, Mann-Whitney U test, Chi-squared test, and Kruskal-Wallis test in biostatistics. *BioData Mining*, 18(1), 56. <https://doi.org/10.1186/s13040-025-00465-6>
- Chiodi, C., Zardinoni, G., Stevanato, P., Giagnoni, L., Carletti, P., Oustrière, N., ... & Renella, G. (2025). Organic amendments influence soil properties, soil microbial diversity, and winter barley traits in a five-year field trial with contaminated soils at a former wood preservation site. *Chemical and Biological Technologies in Agriculture*, 12(1), 87. <https://doi.org/10.1186/s40538-025-00810-1>
- Chitraningrum, N., Gunawan, F., Farma, R., Subyakto, S., Subhan, A., Fudholi, A., ... & Munte, W. N. (2025). Nitrogen-doped activated carbon derived from oil palm empty fruit bunch (OPEFB) for sustainable lithium-ion battery. *Biomass Conversion and Biorefinery*, 15(9), 13845–13860. <https://doi.org/10.1007/s13399-024-06220-x>
- De la Peña, M., Ruiz-Romero, R., & Romero, H. M. (2023). Nitrogen use efficiency in oil palm seedlings: Unraveling the untapped potential of elevated external ammonium supply. *Plants*, 12(15), 2819. <https://doi.org/10.3390/plants12152819>
- Demo, A. H., & Asefa Bogale, G. (2024). Enhancing crop yield and conserving soil moisture through mulching practices in dryland agriculture. *Frontiers in Agronomy*, 6, 1361697. <https://doi.org/10.3389/fagro.2024.1361697>
- Edyson, Murgianto, F., & Ardiyanto, A. (2022). Impact of palm oil by-products application to earthworm populations and soil chemical in oil palm plantation. *IOP Conference Series: Earth and Environmental Science*, 995(1), 012062. <https://doi.org/10.1088/1755-1315/995/1/012062>
- Fang, J., Tao, Y., Liu, J., Lyu, T., Yang, X., Ma, S., ... & Zhang, H. (2023). Effects of emergent plants on soil carbon-fixation and denitrification processes in freshwater and brackish wetlands in a watershed in northern China. *Geoderma*, 430, 116311. <https://doi.org/10.1016/j.geoderma.2022.116311>
- Fedenko, J., D'Amore, D., Spinola, D., Portes, R., Dere, A., & Lybrand, R. A. (2024). Spodosol development and soil organic carbon distribution along a lithosequence in perhumid coastal temperate rainforest. *Soil Science Society of America Journal*, 88(5), 1509–1528. <https://doi.org/10.1002/saj2.20695>
- Ferro-Vázquez, C., Nóvoa-Muñoz, J. C., Klaminder, J., Gómez-Armesto, A., & Martínez-Cortizas, A. (2020). Comparing podzolization under different bioclimatic conditions. *Geoderma*, 377, 114581. <https://doi.org/10.1016/j.geoderma.2020.114581>
- Formaglio, G., Veldkamp, E., Duan, X., Tjoa, A., & Corre, M. D. (2020). Herbicide weed control increases nutrient leaching compared to mechanical weeding in a large-scale oil palm plantation. *Biogeosciences*, 17(21), 5243–5262. <https://doi.org/10.5194/bg-17-5243-2020>
- Guo, H., He, Q., Meng, Q., Min, Z., Chen, P., Li, Y., ... & Xu, J. (2025). The aggregate structure and organic carbon mineralization in forest soils along an elevation gradient in the Sygera Mountains of the southeastern Tibetan Plateau. *Forests*, 16(2), 298. <https://doi.org/10.3390/f16020298>
- Guo, Y., Zhang, X., Zhang, Y., Wu, D., McLaughlin, N., Zhang, S., ... & Liang, A. (2019). Temporal variation of earthworm impacts on soil organic carbon under different tillage systems. *International Journal of Environmental Research and Public Health*, 16(11), 1908. <https://doi.org/10.3390/ijerph16111908>

- Gutierrez Al-Khudhairi, S., Howells, T. R., Bin Sailim, A., McClean, C. J., Senior, M. J. M., Azmi, R., ... & Hill, J. K. (2023). Sustainable management practices do not reduce oil palm yields on smallholder farms on Borneo. *Agroecology and Sustainable Food Systems*, 47(1), 3–24. <https://doi.org/10.1080/21683565.2022.2131691>
- Hallam, J., Berdeni, D., Grayson, R., Guest, E. J., Holden, J., Lappage, M. G., ... & Hodson, M. E. (2020). Effect of earthworms on soil physico-hydraulic and chemical properties, herbage production, and wheat growth on arable land converted to ley. *Science of The Total Environment*, 713, 136491. <https://doi.org/10.1016/j.scitotenv.2019.136491>
- Hau, L. J., Shamsuddin, R., May, A. K. A., Saenong, A., Lazim, A. M., Narasimha, M., & Low, A. (2020). Mixed composting of palm oil empty fruit bunch (EFB) and palm oil mill effluent (POME) with various organics: An analysis on final macronutrient content and physical properties. *Waste and Biomass Valorization*, 11(10), 5539–5548. <https://doi.org/10.1007/s12649-020-00993-8>
- Hu, Y., & Zhu, B. (2025). Effects of organic amendments with different C/N ratios and application rates on N mineralization–immobilization in calcareous cropland soil. *Agronomy*, 15(12), 2795. <https://doi.org/10.3390/agronomy15122795>
- Huang, Y. C., Chen, Z. S., Hsu, J. Y., Chiu, C. M., & Hseu, Z. Y. (2022). Differentiation of fine-textured podzolic soils controlled by climate and landscape in Taiwan. *Geoderma*, 428, 116155. <https://doi.org/10.1016/j.geoderma.2022.116155>
- Iordache, M. (2023). Chemical composition of earthworm casts as a tool in understanding the earthworm contribution to ecosystem sustainability - a review. *Plant, Soil and Environment*, 69(6), 247–268. <https://doi.org/10.17221/461/2022-PSE>
- James, S. W., Bartz, M. L. C., Stanton, D. W. G., Conrado, A. C., Dupont, L., Taheri, S., ... & Brown, G. G. (2019). A neotype for *Pontoscolex corethrurus* (Müller, 1857) (Clitellata). *Zootaxa*, 4545(1), 124–132. <https://doi.org/10.11646/zootaxa.4545.1.7>
- Januari, A. D., Utomo, S. W., & Agustina, H. (2020). Estimation and potential of palm oil empty fruit bunches based on crude palm oil forecasting in Indonesia. *E3S Web of Conferences*, 211, 05003. <https://doi.org/10.1051/e3sconf/202021105003>
- Kamireddy, M., Behera, S. K., & Kancherla, S. (2023). Establishing critical leaf nutrient concentrations and identification of yield limiting nutrients for precise nutrient prescriptions of oil palm (*Elaeis guineensis* Jacq) plantations. *Agriculture*, 13(2), 453. <https://doi.org/10.3390/agriculture13020453>
- Kaupper, T., Hetz, S., Kolb, S., Yoon, S., Horn, M. A., & Ho, A. (2020). Deforestation for oil palm: Impact on microbially mediated methane and nitrous oxide emissions, and soil bacterial communities. *Biology and Fertility of Soils*, 56(3), 287–298. <https://doi.org/10.1007/s00374-019-01421-3>
- Khoramizadeh, A., Jourgholami, M., Jafari, M., Venanzi, R., Tavankar, F., & Picchio, R. (2021). Soil restoration through the application of organic mulch following skidding operations causing vehicle induced compaction in the Hyrcanian Forests, Northern Iran. *Land*, 10(10), 1060. <https://doi.org/10.3390/land10101060>
- Kuligiewicz, A., Szymański, W., Bindeman, I., Kowalik-Hyla, M., Ciesielska, Z., Ziemiański, P. P., & Róžański, K. (2024). Links between environmental conditions and pedogenic phyllosilicates in Podzols in the Tatra Mountains (Poland) – I. Mineralogy. *Geoderma*, 449, 116981. <https://doi.org/10.1016/j.geoderma.2024.116981>
- Lew, J. H., May, A. K. A., Shamsuddin, M. R., Aqsha, Lazim, A. M., & Narasimba, M. M. (2020). Vermicomposting of palm oil empty fruit bunch (EFB) based fertilizer with various organics additives. *IOP Conference Series: Materials Science and Engineering*, 736(5), 052014. <https://doi.org/10.1088/1757-899X/736/5/052014>
- Li, K., Liu, D. N., Li, L. Y., Gao, Y., Gao, W. J., Chen, B. W., ... & Yao, Y. (2025). Effects of nitrogen application amount on nitrogen distribution and photosynthesis in tea leaves. *Frontiers in Plant Science*, 16, 1575317. <https://doi.org/10.3389/fpls.2025.1575317>
- Lopes, I., de Lima, J. L. M. P., Montenegro, A. A. A., & Carvalho, A. A. de. (2025). Assessment of water retention and absorption of organic mulch under simulated rainfall for soil and water conservation. *Soil Systems*, 9(1), 4.

- <https://doi.org/10.3390/soilsystems9010004>
- Ma, L., Song, D., Liu, M., Li, Y., & Li, Y. (2022). Effects of earthworm activities on soil nutrients and microbial diversity under different tillage measures. *Soil and Tillage Research*, 222, 105441. <https://doi.org/10.1016/j.still.2022.105441>
- Mahardika, M., Zakiyah, A., Ulfa, S. M., Ilyas, R. A., Hassan, M. Z., Amelia, D., ... & Norrahim, M. N. F. (2024). Recent developments in oil palm empty fruit bunch (OPEFB) fiber composite. *Journal of Natural Fibers*, 21(1), 2309915. <https://doi.org/10.1080/15440478.2024.2309915>
- Mamabolo, E., Pryke, J. S., & Gaigher, R. (2024). Soil macrofauna are important bioindicators of soil quality in agroecosystems under different management. *Ecological Indicators*, 167, 112723. <https://doi.org/10.1016/j.ecolind.2024.112723>
- Martial, N. G., Yao, S., Hamer, U., Zhang, Y., & Zhang, B. (2023). Positive and negative priming effects induced by freshly added mineral-associated oxalic acid in a Mollisol. *Rhizosphere*, 26, 100708. <https://doi.org/10.1016/j.rhisph.2023.100708>
- Maticic, M., Dugan, I., & Bogunovic, I. (2024). Challenges in sustainable agriculture—The role of organic amendments. *Agriculture*, 14(4), 643. <https://doi.org/10.3390/agriculture14040643>
- Mazzilli, S. R., Kemanian, A. R., Ernst, O. R., Jackson, R. B., & Piñeiro, G. (2014). Priming of soil organic carbon decomposition induced by corn compared to soybean crops. *Soil Biology and Biochemistry*, 75, 273–281. <https://doi.org/10.1016/j.soilbio.2014.04.005>
- Nadolny, H., Santos, A., Demetrio, W., Ferreira, T., Maia, L. D. S., Conrado, A. C., ... & Brown, G. (2020). Recommendations for assessing earthworm populations in Brazilian ecosystems. *Pesquisa Agropecuária Brasileira*, 55, e01006. <https://doi.org/10.1590/s1678-3921.pab2020.v55.01006>
- Nelson, D., & Sommers, L. (1982). Total carbon, organic carbon, and organic matter. Methods of soil analysis. Part 2. *Chemical and Microbiological Properties*, 9, 539–579. <https://doi.org/10.2134/agronmonogr9.2.2ed.c29>
- Noirot, L. M., Müller-Stöver, D. S., Wahyuningsih, R., Sørensen, H., Sudarno, Simamora, A., ... & Caliman, J.-P. (2022). Impacts of empty fruit bunch applications on soil organic carbon in an industrial oil palm plantation. *Journal of Environmental Management*, 317, 115373. <https://doi.org/10.1016/j.jenvman.2022.115373>
- Nyasapoh, J. B. A., Opong Danso, E., Arthur, E., Obour, P. B., Amponsah, W., Sabi, E. B., ... & Andersen, M. N. (2026). Oil palm empty fruit bunch amendments improved the structural and functional properties of a tropical Acrisol. *Soil Science Society of America Journal*, 90(1), e70195. <https://doi.org/10.1002/saj2.70195>
- Panta, U., Jernigan, A., & Rashed, A. (2026). Effects of soil biofumigation on non-target springtails (*Collembola*) and earthworms (*Opisthopora*). *Scientific Reports*, 16(1), 3551. <https://doi.org/10.1038/s41598-025-33634-y>
- Petticord, D. F., Uveges, B. T., Boughton, E. H., Strahm, B. D., & Sparks, J. P. (2025). Spatial decoupling of biological and geochemical phosphorus cycling in podzolized soils. *Soil Systems*, 9(4), 115. <https://doi.org/10.3390/soilsystems9040115>
- Phillips, I., Paungfoo-Lonhienne, C., Tahmasbian, I., Hunter, B., Smith, B., Mayer, D., & Redding, M. (2022). Combination of inorganic nitrogen and organic soil amendment improves nitrogen use efficiency while reducing nitrogen runoff. *Nitrogen*, 3(1), 58–73. <https://doi.org/10.3390/nitrogen3010004>
- Potapov, A., Schaefer, I., Jochum, M., Widyastuti, R., Eisenhauer, N., & Scheu, S. (2021). Oil palm and rubber expansion facilitates earthworm invasion in Indonesia. *Biological Invasions*, 23(9), 2783–2795. <https://doi.org/10.1007/s10530-021-02539-y>
- Radics, Z., Dekemati, I., Gyuricza, C., Simon, B., Ibrahim, H., Vinogradov, S., & Birkás, M. (2022). Effects of irrigation and organic mulching on the abundance and biomass of earthworms. *Polish Journal of Environmental Studies*, 31(3), 2811–2821. <https://doi.org/10.15244/pjoes/143258>
- Rahman, N., Giller, K. E., de Neergaard, A., Magid, J., van de Ven, G., & Bruun, T. B. (2021). The effects of management practices on soil organic carbon stocks of oil palm plantations in Sumatra, Indonesia. *Journal*

- of Environmental Management*, 278, 111446. <https://doi.org/10.1016/j.jenvman.2020.111446>
- Ratsiatosika, O., Trap, J., Herinasandratra, V., Razafimbelo, T., Bernard, L., & Blanchart, E. (2024). Earthworms enhance the performance of organic amendments in improving rice growth and nutrition in poor ferralsols. *Soil Biology and Biochemistry*, 195, 109477. <https://doi.org/10.1016/j.soilbio.2024.109477>
- Readyhough, T., Neher, D. A., & Andrews, T. (2021). Organic amendments alter soil hydrology and belowground microbiome of tomato (*Solanum lycopersicum*). *Microorganisms*, 9(8), 1561. <https://doi.org/10.3390/microorganisms9081561>
- Rizinjirabake, F., Tenenbaum, D. E., & Pilesjö, P. (2019). Sources of soil dissolved organic carbon in a mixed agricultural and forested watershed in Rwanda. *CATENA*, 181, 104085. <https://doi.org/10.1016/j.catena.2019.104085>
- Rosa Silva, A. A., da Silva Brito, G. F., Araruna, T., Gambetta, R., & Machado, F. (2026). Thermochemical conversion of oil palm empty fruit bunch into fuel gas in a fluidized bed gasifier. *Bioresource Technology*, 440, 133404. <https://doi.org/10.1016/j.biortech.2025.133404>
- Rosenani, A. B., Rabuni, W., Cheah, P., & Noraini, J. (2016). Mass loss and release of nutrient from empty fruit bunch of oil palm applied as mulch to newly transplanted oil palm. *Soil Research*, 54(8), 985–996. <https://doi.org/10.1071/SR15143>
- Rudolf, K., Hennings, N., Dippold, M. A., Edison, E., & Wollni, M. (2021). Improving economic and environmental outcomes in oil palm smallholdings: The relationship between mulching, soil properties and yields. *Agricultural Systems*, 193, 103242. <https://doi.org/10.1016/j.agsy.2021.103242>
- Santos, F. C. D., Albuquerque Filho, M. R. D., Soares, J. R., Jesmin, T., Manirakiza, N., Silveira, M. C. T. D., ... & Bhadha, J. H. (2025). Organic matter: A critical soil health indicator in agrosystems within the Cerrado of Bahia, Brazil and South Florida, USA. *Frontiers in Soil Science*, 5, 1524550. <https://doi.org/10.3389/fsoil.2025.1524550>
- Sari, W. K., & Malik, P. A. (2023). The effects of application of biochar from oil palm empty fruit bunches on chemical properties of ultisols and the growth of cacao seedlings. *Kultivasi*, 22(2), 157–167. <https://doi.org/10.24198/kultivasi.v22i2.46525>
- Shabtai, I. A., Hafner, B. D., Schweizer, S. A., Höschen, C., Possinger, A., Lehmann, J., & Bauerle, T. (2024). Root exudates simultaneously form and disrupt soil organo-mineral associations. *Communications Earth & Environment*, 5(1), 699. <https://doi.org/10.1038/s43247-024-01879-6>
- Shumba, A., Chikowo, R., Thierfelder, C., Corbeels, M., Six, J., & Cardinael, R. (2024). Mulch application as the overarching factor explaining increase in soil organic carbon stocks under conservation agriculture in two 8-year-old experiments in Zimbabwe. *Soil*, 10(1), 151–165. <https://doi.org/10.5194/soil-10-151-2024>
- Souza Junior, A. J. de, Camêlo, D. de L., Arruda, D. L. de, Souza Junior, V. S. de, Rocha, A. T. da, & Corrêa, M. M. (2023). Spodosol formation on sandy ruins in a semi-arid climate in the Catimbau National Park, Northeast Brazil. *CATENA*, 229, 107226. <https://doi.org/10.1016/j.catena.2023.107226>
- Statistics Indonesia. (2025). *Indonesia Permanent Estate Crops Statistics 2024* (Vol. 1). Retrieved from <https://www.bps.go.id/en/publication/2025/08/29/8d2a6ab3510f9828daf73191/statistik-tanaman-perkebunan-tahunan-indonesia-2024-kelapa-sawit-kopi-kakao-karet-teh-dan-komoditas-perkebunan-unggulan.html>
- Supriatna, J., Setiawati, M. R., Sudirja, R., Suherman, C., & Bonneau, X. (2022). Composting for a more sustainable palm oil waste management: A systematic literature review. *The Scientific World Journal*, 2022(1), 5073059. <https://doi.org/10.1155/2022/5073059>
- Syahza, A., & Asmit, B. (2019). Regional economic empowerment through oil palm economic institutional development. *Management of Environmental Quality: An International Journal*, 30(6), 1256–1278. <https://doi.org/10.1108/MEQ-02-2018-0036>
- Taheri, S., Pelosi, C., & Dupont, L. (2018). Harmful or useful? A case study of the exotic peregrine earthworm morphospecies *Pontoscolex corethrurus*. *Soil Biology and*

- Biochemistry*, 116, 277–289. <https://doi.org/10.1016/j.soilbio.2017.10.030>
- Tao, H. H., Snaddon, J. L., Slade, E. M., Henneron, L., Caliman, J. P., & Willis, K. J. (2018). Application of oil palm empty fruit bunch effects on soil biota and functions: A case study in Sumatra, Indonesia. *Agriculture, Ecosystems & Environment*, 256, 105–113. <https://doi.org/10.1016/j.agee.2017.12.012>
- Tarigan, R. S., Suhardi, Wahyuningsih, R., Pujianto, Wahyuningsih, P. A., Marichal, R., & Caliman, J. P. (2019). The role of oil palm biomass recycling on soil health in oil palm plantations. *IOP Conference Series: Earth and Environmental Science*, 336(1), 012019. <https://doi.org/10.1088/1755-1315/336/1/012019>
- Tecimen, H. B., Gorres, J. H., & Melnichuk, R. D. S. (2021). Effect of *Lumbricus rubellus* and *Amyntas agrestis* earthworms on soil biogeochemistry at the aggregate scale in Northern Hardwood Forests. *Journal of Sustainable Forestry*, 40(1), 83–98. <https://doi.org/10.1080/10549811.2020.1744452>
- Tianle, L., Quanhua, Z., Shuhan, J., Yu, L., & Xueliang, G. (2026). Enhanced soil organic matter estimation using spectral feature selection with empirical mode decomposition and pearson correlation coefficients. *Infrared Physics & Technology*, 152, 106084. <https://doi.org/10.1016/j.infrared.2025.106084>
- Traoré, S., Guébré, D., Hien, E., Traoré, M., Lee, N., Lorenz, N., & Dick, R. P. (2022). Nutrient cycling and microbial responses to termite and earthworm activity in soils amended with woody residues in the Sudano-Sahel. *European Journal of Soil Biology*, 109, 103381. <https://doi.org/10.1016/j.ejsobi.2021.103381>
- Utomo, Y. M. P., Risnawati, R., Rohimsyah, F. M., Tominaga, M., Kurniawan, F., Astuti, W., ... & Triana, Y. (2024). Oil palm empty fruit bunch (OPEFB) activated carbon as promising electrode materials for battery. *Solid State Phenomena*, 368, 61–76. <https://doi.org/10.4028/p-fDu5JK>
- van Rijssel, S. Q., Kuipers, E., Mason-Jones, K., Koorneef, G. J., van der Putten, W. H., & Veen, G. F. (2025). Impact of soil inoculation on crop residue breakdown and carbon and nitrogen cycling in organically and conventionally managed agricultural soils. *Applied Soil Ecology*, 205, 105760. <https://doi.org/10.1016/j.apsoil.2024.105760>
- Vanella, D., Guarrera, S., Ferlito, F., Longo-Minnolo, G., Milani, M., Pappalardo, G., ... & Consoli, S. (2025). Effects of organic mulching and regulated deficit irrigation on crop water status, soil and yield features in an orange orchard under Mediterranean climate. *Science of The Total Environment*, 958, 177528. <https://doi.org/10.1016/j.scitotenv.2024.177528>
- Vršič, S., Breznik, M., Pulko, B., & Rodrigo-Comino, J. (2021). Earthworm abundance changes depending on soil management practices in Slovenian Vineyards. *Agronomy*, 11(6), 1241. <https://doi.org/10.3390/agronomy11061241>
- Wang, J., Yang, X., Huang, S., Wu, L., Cai, Z., & Xu, M. (2024). Long-term combined application of organic and inorganic fertilizers increases crop yield sustainability by improving soil fertility in maize-wheat cropping systems. *Journal of Integrative Agriculture*, 24(1), 290–305. <https://doi.org/10.1016/j.jia.2024.07.003>
- Wang, M., Wang, R., Jiang, G., Li, Y., Liu, C., Yang, J., ... & Liu, S. (2025). Soil organic carbon and ecosystem multifunctionality are enhanced by subsoiling in fluvo-aquic soil of North China Plain. *Frontiers in Plant Science*, 16, 1559653. <https://doi.org/10.3389/fpls.2025.1559653>
- Ward, E. B., & Bradford, M. A. (2025). Plant nitrogen demand, not soil carbon availability, decouples net mineralization and nitrification following forest canopy disturbances. *Biogeochemistry*, 168(4), 59. <https://doi.org/10.1007/s10533-025-01251-6>
- Wierzbowska, J., Sienkiewicz, S., & Załuski, D. (2021). Nitrogen fractions in soil fertilized with waste organic materials. *Agronomy*, 11(8), 1474. <https://doi.org/10.3390/agronomy11081474>
- Xu, J., Li, Y., & Li, L. (2025). A comprehensive review of the effects of organic amendments on soil health and fertility: Mechanisms, greenhouse gas emissions, and implications for sustainable agriculture. *Agronomy*, 15(12), 2705. <https://doi.org/10.3390/agronomy15122705>

- Yadegari, M., Shamshiri, R. R., Mohamed Shariff, A. R., Balasundram, S. K., & Mahns, B. (2020). Using SPOT-7 for nitrogen fertilizer management in oil palm. *Agriculture*, *10*(4), 133. <https://doi.org/10.3390/agriculture10040133>
- Yaseen, A., & Abdulqadir, M. K. (2024). Mulching strategies and the significance of mulching in improving soil fertility and soil physical properties: A review. *Acta Agriculturae Slovenica*, *120*(4), 1–15. <https://doi.org/10.14720/aas.2024.120.4.18807>
- Zhang, L., Liu, J., Yin, R., Xu, Z., You, C., Li, H., ... & Tan, B. (2023). Soil fauna accelerated litter C and N release by improving litter quality across an elevational gradient. *Ecological Processes*, *12*(1), 47. <https://doi.org/10.1186/s13717-023-00459-4>
- Zhelezova, A., Sorrentino, G., Otim, G. I., & Rocchi, I. (2025). Soil is alive – How does soil biota influence soil mechanical properties? A perspective review. *Biogeotechnics*, 100175. <https://doi.org/10.1016/j.bgtech.2025.100175>