



Evaluation of phosphate sorption capacity and external phosphorus requirement of some agricultural soils of the southwestern Ethiopian highlands

Berhanu Dinssa¹, Eyasu Elias²

¹ Department of Plant Science, Ambo College of Agriculture, Ambo University, Ethiopia

² Centre of Environmental Sciences, College of Natural Sciences, Addis Ababa University, Ethiopia

ARTICLE INFO

Keywords:

External P requirement
Freundlich equation
Phosphorous
P-sorption capacity

Article history

Submitted: 2021-05-20

Accepted: 2021-11-09

Available online: 2021-12-26

Published regularly: December 2021

* Corresponding Author

Email address:

berhanudj@gmail.com

ABSTRACT

One of the most soil fertility management problems for crop production on acidic soils of the Ethiopian highlands is phosphorus fixation. The research was executed to assess the P-sorption capacity and to determine the external P requirement of different acidic soils in the Southwestern highlands of Ethiopia. Phosphorus sorption capacity (Kf) and its relation with selected soil characteristics were assessed for some major agricultural soils in the Ethiopian highlands to answer the questions, 'What are the amount of P-sorption capacity and external P requirement of Nitisols, Luvisols, Alisols, and Andosols in Ethiopia?'. Twelve surface soil samples (at depth of 0-30 cm) were gathered and the P-sorption capacity was estimated. Phosphorus-sorption data were obtained by equilibrating 1 g of the 12 soil samples with 25 ml of KH_2PO_4 in 0.01 M CaCl_2 , having 0, 30, 60, 90, 120, 150, 180, 210, 240, 270, 300, and 330 mg P L^{-1} for 24 hours. The data were adjusted to the Freundlich adsorption model and the relationship among P-sorption and soil characteristics was established by correlation analysis. Clay content and exchangeable acidity, organic matter, Al_2O_3 , and Fe_2O_3 oxides have affected phosphorus-sorption at a significance level of ($P < 0.05$). Alisols had the highest Kf value (413 mg kg^{-1}) but Nitisols had the lowest Kf (280 mg kg^{-1}). The external phosphorus fertilizer requirement of the soils was in the order of 25, 30, 32, and 26 mg P kg^{-1} for Nitisols, Luvisols, Alisols, and Andosols sequentially. The Kf varies among different soil types of the study area. The magnitude of the soil's Kf was affected by the pH of the soil, soil OM content, and oxides of Fe and Al. Therefore, knowledge of the soils' P retention capacity is highly crucial to determine the correct rate of P fertilizer for crop production.

How to Cite: Dinssa, B. & Elias, E. (2021). Evaluation of phosphate sorption capacity and external phosphorus requirement of some agricultural soils of the southwestern Ethiopian highlands. *Sains Tanah Journal of Soil Science and Agroclimatology*, 18(2): 136-142. <https://dx.doi.org/10.20961/stjssa.v18i2.51325>

1. INTRODUCTION

Ethiopia has a diversity of topography, climate, vegetation, and parent materials, as a soil-forming factor, leading to the formation of various soil types (Elias, 2016). Nitisols and Luvisols are among the most extensive soil types covering about 31 and 11% of the arable landscapes of the Ethiopian highlands, whereas the highly productive Andosols are less extensively distributed (< 1%) (Elias et al., 2019). Alisols are particularly important in high rainfall upland farming systems of the Southwestern highlands of Ethiopia. Regardless of their area extent and agricultural importance, the phosphorus limitation of these soils has been constraining agricultural productivity in the Ethiopian highlands. Preliminary reports about the nutrient content of the Ethiopian soils suggest that 99% of the Ethiopian highland

soils are phosphorus-deficient by taking 30 ppm of available phosphorus as a critical level (Karlton et al., 2013). Studies on the nutrient status of the Ethiopian highland soils suggest that phosphorus deficiency is a major problem that negatively affects crop production (Elias et al., 2019; Melese et al., 2015).

A widespread problem of limited P-availability in soils might be due to severe P-fixation. Phosphate fixation is a process by which phosphate ions are retained on the active sites of the soil colloidal surfaces thus rendering it unavailable for plant uptake (Tamungang et al., 2014). Phosphorus fixation is a common problem in tropical rainfed upland farming systems (Hanyabui et al., 2020). The high P-retention by adsorption or precipitation reactions in soils results in low

P-fertilizer use efficiency (15–30%) (Maluf et al., 2018). The estimated 80% of the applied phosphate is not available for crop uptake as the result of high phosphorus fixation in acid tropical soils (Kisinyo et al., 2015). Most inherent characteristics of the soil that affect the degree of P-fixation include soil clay content, soil reaction (pH), OM content, hydrated oxides of Fe and Al, and the concentration of exchangeable aluminum and hydrogen (exch. Al^{3+} and H^+) that varies with soil type (Wang et al., 2017). Likewise, Jiang et al. (2015) stated that the oxides of Fe and Al are the major adsorbents for phosphate ions in acidic tropical soils. When the clay fraction is composed of 1:1 minerals, iron, and aluminum oxides and hydroxides, most of the P is adsorbed on the soil colloidal surfaces. The phosphorus sorption capacity (K_f) of a soil refers to the potential of the soil to bind phosphorus applied to it and is a principal factor regulating P concentration in the soil solution and available phosphorus (Gonzalez-Rodriguez & Fernandez-Marcos, 2018). The external P requirement (EPR), also described as the standard P requirement (SPR), is the amount of P which must be applied to the soil to sustain a P level in the soil at 0.2 mg P L^{-1} that meets the requirements in P for different crops (Wolde & Haile, 2015).

Determination of the K_f and EPR_f of various types of soils supports sustainable agriculture through site-specific P fertilizer recommendations that avoid overdose P application that led to adverse impacts on the economy and the environment. Underdoes application of P negatively impacts the growth potential of the crops and affects the living standard of the society through food self-insufficiency. Similarly, Thuy et al. (2020) stated that optimizing P fertilizer input could maximize the P fertilizers use efficiency that will decrease fertilizer costs, improve income, and lower the negative impacts on the soil for sustainable agriculture.

Different mathematical models and sorption isotherms have been formulated to explore the soils P sorption capacity (K_f); P-sorption energy ($1/n$), buffering capacity of soil versus P concentration gradient in solution, and the state of equilibrium between P in solution and the solid phases and their relation (Muindi et al., 2015). Freundlich model is the

most commonly applicable model to determine P-sorption characteristics of soils and estimate EPR_f (Afsar et al., 2012). This model can differentiate soils according to their capacity to absorb P from soil solutions. Although studies have indicated the widespread P-sorption problem in Ethiopian soils (Wolde & Haile, 2015). Information on P-sorption disaggregated by soil type is rather limited in the Southwestern highlands of Ethiopia. In recent years, Ethiopia is in the process of transforming its soil fertility sector by moving away from blanket fertilizer recommendation to soil-specific management of fertilizer (Elias et al., 2019). To this effect, to alleviate the P-sorption problem through soil-specific management (e.g., liming and targeted P-fertilization), it is necessary to estimate the degree of P-fixation by dominant soil types. Hence, this research intended to evaluate the P-sorption capacity and external P requirement of some acidic soils (Nitisols, Luvisols, Alisols, and Andosols) in the Southwestern highlands of Ethiopia.

2. MATERIAL AND METHODS

2.1. Description of study sites

The experiment was carried out on four dominant agricultural soils based on FAO classification (Nitisols, Luvisols, Alisols, and Andosols) in the intervention sites of the project; Capacity building for scaling up of evidence-based best practices in agricultural production in Ethiopia (CASCAPE) intervention woredas of Oromiya region, (Fig.1). The district is designated by flat plains, high mountains, and hilly topography, geological feature of the area is characterized by tertiary sediments of the Cenozoic era on the plain and basalt and volcanic rocks in the High Mountain and hilly ridges. The rainfall data obtained from the nearest weather stations (Bako and Jimma Agricultural Research Centers) reveals that the rainy season covers April to November and maximum rain is received in June, July, and August. The long-term (2009- 2018) mean annual rainfall is 1267-1730 mm. It has a warm humid climate with average air temperatures of 19 to 21 °C and the altitudinal range of the sites is 1650 to 2800 m asl.




Figure 1. Location map of the study area

2.2. Soil sampling and laboratory procedures

About ten representative sub-samples were taken from 0-30 cm depth randomly in a zig-zag manner from different spots of the sampling unit and one composite sample was formed (1 kg each). The sub-samples were thoroughly mixed together on a flat plastic sheet and 12 composite soil samples (3 samples from every type of soil) were collected in clean new plastic bags and transported to HortiCoops soil fertility laboratory. In the laboratory, the samples were dried on the air, crushed, sieved, and passed through a 2 mm mesh, to determine the soil physicochemical properties and P sorption capacity.

The soil particle size distribution was determined by the hydrometer method (Beretta et al., 2014). Soil pH (H₂O) was measured by a pH meter as described by (FAO, 2021). OM (%) was determined by the wet digestion method (FAO, 2020). Exch. Al³⁺ and H (cmol₍₊₎kg⁻¹) were described by leaching with KCl followed by titration (Lestari et al., 2016). Determination of Fe₂O₃ and Al₂O₃ was following the X-ray fluorescence (XRF) spectrometer (Towett et al., 2013).

Phosphorus sorption characteristics of the soil were analyzed by using three replicates of 1.0 g, dry soil samples that weighed into 50 ml centrifuge tubes and suspended in 25 ml of 0.01 M CaCl₂ as supporting electrolytes that have 0, 30, 60, 90, 120, 150, 180, 210, 240, 270, 300 and 330 mg P L⁻¹ as KH₂PO₄ with increasing rate of 30 mg P L⁻¹. Three drops of toluene were applied to all containers to stop the microbial action. The tubes were then closed and shaken on an end-to-end shaker for 24 hours at a temperature of 25 ± 1 °C at 150 oscillations per minute. After equilibrium, the soil suspensions were centrifuged at about 3000 rpm for 10 minutes, then filtered through Whatman No. 42 to obtain a clear solution. The phosphorus in the supernatant was then analyzed following the Olsen procedure. The quantity of P absorbed was calculated as the difference between the amount of P added and that of left in solution by Equation 1.

$$P_{\text{absorbed}} = \frac{(C_0 - C_f)V}{\text{Weight of soil (kg)}} \quad [1]$$

Where: C₀ = initial concentration of P (mg L⁻¹), C_f = final concentration of P (mg L⁻¹) and V = volume of solution (L). The soil P sorption data were adjusted to the linearized forms of Freundlich (Equation 2).

$$X = KC^{1/n} \quad [2]$$

The logarithm of Eq. [2], changes into log X = log K_f + 1/n log C, the linear equation form 2 has been formulated by plotting Log C against Log X, X (mg kg⁻¹) = the concentration of P retained per unit mass of soil, C (mg L⁻¹) = the equilibrium concentration of P, 1/n (L kg⁻¹) represents the slope (the sorption energy) and Log K_f (mg kg⁻¹) represents the sorption capacity in mg kg⁻¹ (Dari et al., 2015). The external phosphorus requirement (EPR_f) is calculated based on Freundlich models by using the required P quantity (0.2 mg P L⁻¹) in the linear form of Freundlich equations (Getie et al., 2021). The P-buffering capacity (PBC) of a soil is its capacity to resist change in the P concentration of the solution phase (Shirvani et al., 2005). It was computed from the Freundlich adsorption isotherms at 0.2 mg L⁻¹ of solution P as PBC (Eq. 3).

$$PBC = K_f * n * C^{n-1} \quad [3]$$

Whereas: PBC (L kg⁻¹) = Phosphorus buffering Capacity; K_f (mg kg⁻¹) = P sorption capacity; 1/n (kg L⁻¹) = sorption energy; C (mg L⁻¹) = the equilibrium concentration of P. The citrate bicarbonate dithionite (CBD) procedure was used for the determination of total free iron oxides in soils (Fan et al., 2016).

2.3 Statistical analysis

The measured soil parameters were arranged in a Completely Randomized Design (CRD) with three replicates and subjected to analysis of variance (ANOVA) using the Statistical Analysis System (SAS) software version 9.4 (SAS Institute Inc, 2019). The Least Significance Difference test (LSD) at the 5% probability level was used for the mean comparison when the ANOVA showed significant differences. Also, a simple regression and correlation coefficient analysis was executed to assess the degrees and directions of the relationship between K_f and the various soil parameters.

3. RESULTS

3.1. Phosphorus sorption capacity of soils

Table 1 shows highly significant (P < 0.01) variation among soils types for P-sorption capacity (K_f), level of P-sorption energy (1/n) (P < 0.05), and phosphorus buffering capacity (PBC_f) (P < 0.01) but no significant difference in the external phosphorus requirement (Table 1).

Table 1. P-sorption capacity (K_f), Slope (n), EPR_f and PBC_f values after 24 hours of equilibration

Soil Type (RSG)	K _f (mg kg ⁻¹)	1/n (L kg ⁻¹)	EPR _f (mg kg ⁻¹)	PBC _f (L kg ⁻¹)	R ²
Nitisols	280 ^b	1.40 ^b	25	316 ^b	0.95 ^b
Luvisols	336 ^{ab}	1.48 ^{ab}	30	395 ^{ab}	0.97 ^a
Alisols	413 ^a	1.67 ^a	32	471 ^a	0.98 ^a
Andosols	299 ^b	1.43 ^b	26	346 ^b	0.96 ^{ab}
Mean	332	1.49	28	382	0.97
LSD%	0.15	66.07	37.41	80.27	0.003
SE	32.79	0.08	5.0	37.76	0.007
F	9.70	7.46	1.52	9.70	7.59
P	0.01	0.02	0.30	0.01	0.018

Note: In the same column, means indicated by different letters are significantly varied at p < 0.05; K_f = P sorption capacity; 1/n = sorption energy; EPR_f = Freundlich External P Requirement; PBC_f = P buffering capacity; RSG = Reference soil group.




Figure 2. P sorption Isotherm for Nitisols (a), Luvisols (b), Alisols (c) and Andosols (d)

The sequence of soils based on their K_f was Alisols > Luvisols > Andosols > Nitisols. Alisols had the highest K_f (413 mg kg^{-1}) and $1/n$ (1.67 kg L^{-1}) but Nitisols had the lowest K_f (280 mg kg^{-1}) and $1/n$ (1.40 kg L^{-1}). The linear form of the Freundlich equation formed by taking $\log C$ versus $\log X$ is displayed in Figure 2. It confirmed a satisfactory agreement with the Freundlich equation ($R^2 = 0.96\text{--}0.98$) for the studied soils. The P sorption isotherms were formed by plotting the amount of P adsorbed on the soil surface (Nitisols, Luvisols, Alisols, and Andosols) against the solution P concentration.

3.2. The Soil External Phosphorus Requirement (EPR_f) and Phosphorus Buffering Capacity (PBC)

The actual quantities of P required by Nitisols, Luvisols, Alisols, and Andosols were 25, 30, 32, and 26 (mg kg^{-1}) respectively for optimum crop production at the study area (Table 1). The highest (32 mg kg^{-1}) and the least (25 mg kg^{-1}) values of EPR_f were obtained for Alisols and Nitisols respectively. The buffering capacity of soils was computed from the Freundlich adsorption isotherms at 0.2 mg P L^{-1} of solution in the 24 hours of equilibration were 316, 396, 471, and 346 for Nitisols, Luvisols, Alisols, and Andosols respectively (Table 1).

Table 2 shows there is variation in soil characteristics and K_f among the soils. Accordingly, the mean clay content was significantly varied ($p < 0.05$) among various soil types, and the highest value (54.67%) was observed under Alisols while the lowest value (36.0) was under Andosols. Even though statistically no significant difference in pH between the soils, Andosols had the highest (6.30) and Alisols had the lowest (4.80) pH. There was a strong and significant ($p < 0.001$) difference in mean OM content between the soils. The highest

value (7.51%) and the lowest value (3.36%) of OM were observed under Andosols and Alisols respectively. Significant ($p < 0.05$) differences were observed in exchangeable aluminum (exch. Al^{3+}) and exchangeable acidity (exch. Ac) of the soils under this study. Alisols had the highest exch. Al^{3+} ($1.18 \text{ cmol}_{(+)}\text{kg}^{-1}$) and exch. Ac ($1.67 \text{ cmol}_{(+)}\text{kg}^{-1}$) but Nitisols had the lowest (null) exch. Al^{3+} and exch. Ac ($0.52 \text{ cmol}_{(+)}\text{kg}^{-1}$). The soils were strongly and significantly ($p < 0.001$) differs in both Al_2O_3 (%) and Fe_2O_3 (%) content. Nitisols had the highest mean Al_2O_3 (26.68%) and Fe_2O_3 (16%) while Andosols had the lowest mean Al_2O_3 (14.05%) and Fe_2O_3 (9.24%). A significant ($p < 0.05$) variation was observed among the soils in K_f (413 mg kg^{-1}). Accordingly, Alisols had the highest and Nitisols had the lowest K_f (280 mg kg^{-1}) values.

3.3. Correlation between soil properties and P-sorption capacity

The relationship between the K_f and selected soil parameter is displayed in Table 3. Statistically, a positive and significant ($P \leq 0.05$) relationship existed between exch. Al^{3+} and K_f . The K_f had a positive and non-significant ($P > 0.05$) relationship with clay content, exch. H^+ , exch. Ac and aluminum oxides (Al_2O_3) and iron oxides (Fe_2O_3). In addition, the K_f value had negative, non-significant ($p > 0.05$) relationships with pH (H_2O) and OM content of the soil. The sorption energy ($1/n$) had a positive non-significant ($P > 0.05$) relation with clay content while it had strongly significant positive relationships with exch. Al^{3+} at ($P < 0.001$) level of significance and for P sorption capacity (K_f) at 5% level of significance. Moreover, the equilibrium concentration of phosphorus, the slope (sorption energy) had non-significant ($P > 0.05$) relationships with soil pH and soil OM content.

Table 2. Variability in soil properties and P-sorption capacity (K_f) for different soil types in Ethiopian highlands

Soil type	Clay (%)	pH-H ₂ O	OM (%)	Exch. Al ³⁺ (cmol kg ⁻¹)	Exch. H ⁺ (cmol kg ⁻¹)	Exch. Ac (cmol kg ⁻¹)	Al ₂ O ₃ (%)	Fe ₂ O ₃ (%)	K _f (mg kg ⁻¹)
Nitisols	51.7 ^a	5.71	3.8 ^b	0.0 ^b	0.52	0.52 ^{ab}	26.7 ^a	16. ^a	280 ^b
Luvissols	48.7 ^{ab}	5.73	3.83 ^b	0.26 ^{ab}	0.35	0.61 ^{ab}	23.9 ^{ab}	15.9 ^a	336 ^{ab}
Alisols	54.7 ^a	4.8	3.36 ^b	1.18 ^a	0.49	1.67 ^a	22.6 ^b	13.3 ^b	413 ^a
Andosols	36 ^b	6.3	7.51 ^a	0.13 ^b	0.11	0.24 ^b	14.1 ^c	9.24 ^c	299 ^b
Mean	47.8	5.63	4.62	0.39	0.37	0.76	21.8	13.6	332
LSD%	15.42	1.97	1.87	1.02	0.87	1.19	3.74	1.67	0.15
SE	5.45	0.69	0.66	0.36	0.30	0.42	1.30	0.59	32.79
F-value	6.79	2.34	25.52	6.50	1.09	6.63	50.8	86.6	9.70
P-value	0.02	0.17	0.008	0.02	0.42	0.02	0.001	<0.001	0.01
Sign.	0.05	Ns	0.001	0.05	Ns	0.05	0.001	0.001	0.05

Notes: In the same column, means indicated by different letters are significantly varied at $p < 0.05$, ns = non-significant ($P > 0.05$), OM = organic matter, Exch Al³⁺ = Exchangeable aluminum, Exch H⁺ = Exchangeable hydrogen, Exch Ac = Exchangeable acidity, Al₂O₃ = aluminum oxide, Fe₂O₃ = iron oxide.

4. DISCUSSION

The variation in the K_f content of the soils could be attributed to the difference in the soil pH, exch. Al³⁺, clay, and OM in the soils and the percentage of Al₂O₃ and Fe₂O₃ in the soil (Table 2). Very high clay content (54.7%), strongly acidic soil reaction (pH 4.8), high level of exch. Al³⁺ (1.18 mg kg⁻¹) in the soil solution and moderate percentage of OM (3.36%) content in Alisols resulted in the highest K_f (413 mg kg⁻¹) and EPR_f (32.33 mg kg⁻¹) while, moderately acidic soil reaction (pH 5.7), high OM content (3.8%) and nil exch. Al³⁺ in Nitisols attributed to the lowest K_f (280 mg kg⁻¹) and EPR_f (25 mg kg⁻¹). Likewise, the K_f and EPR_f of Luvisols and Andosols were varied based on these soil properties. This finding is similar to the results of (Wang et al., 2017). The K_f values of the soils of this study were the intermediate values (280 - 413 mg P kg⁻¹) detected from (Melese et al., 2015) for the soils of Farta district, in the Northwestern Highlands of Ethiopia which varies from 80 to 259 mg P kg⁻¹ and those K_f value stated by Wolde and Haile (2015) in soils from Bule District, in the Southern Ethiopia, which varies from 479 to 487 mg P kg⁻¹. According to the rating established by Buresh et al. (1997) Nitisols, Luvisols, and Andosols were classified under moderate Phosphorus sorbing (280-316 mg P kg⁻¹) soils. Alisols with K_f values of 413 mg P kg⁻¹) were grouped under high P sorbing soils (Table 2).

The Soil External Phosphorus Requirement (EPR_f), Alisols with the highest PBC (471 L kg⁻¹) had the highest capacity to restore a unit change in soil solution P and Keep a productive solution P concentration and make lower the amount of EPR_f. Because of this, the external phosphorous requirement was not significantly differed among the soil types. The observed EPR_f in the soils is more than double of the previously recommended amounts (20 kg P ha⁻¹) for Alisols and Andosols, and 30 kg P for Nitisols and Luvisols. The result of this study revealed the need for twice phosphorus fertilizer application compared to the previously used phosphorus fertilizer of 30 kg P ha⁻¹ for maize production at Bako and 20 kg P ha⁻¹ at Omonada Districts of Oromiya region that supply only 15-10 mg P kg⁻¹, which is half of the soils' EPR_f of 32-25 mg P kg⁻¹. According to this finding, 50, 60, 64, and 52 kg P ha⁻¹ for Nitisols, Luvisols, Alisols, and Andosols respectively were

recommended for the study areas. Similarly, the Ethio SIS team, Karltun et al. (2013) suggested an initial critical value of 30 mg P kg⁻¹ soil (60 kg P ha⁻¹) for soil fertility mapping and P fertilizer recommendations in Ethiopia was comparable to the finding of this study.

Correlation between Soil Properties and P-Sorption Capacity, The reason for clay particles positively correlated with soils' sorption capacity (K_f) was due to, the fact that finer smaller soil particles (clays) are more reactive and have greater P-retention capacity due to the larger surface area compared to larger soil particles (Jiang et al., 2015). Wehrauch and Opp (2018) also stated that most of the time clay minerals have a more net negative charge than chelating metal cations than sorb Phosphate. The soil pH was negatively correlated with K_f . As the pH increased, K_f decreased due to the increase in electrostatic repulsion created by the increase in negative surface charges.

Table 3. The correlation coefficient between soil properties and P-sorption capacity (K_f)

Selected Soil properties	Correlation coefficient (r)	p-value	Significance
Clay content vs K _f	0.55	0.45	Ns
K _f vs pH-H ₂ O	-0.82	0.18	Ns
K _f vs OM	-0.54	0.46	Ns
K _f vs Exch. Al ³⁺	0.96*	0.04	*
K _f vs Exch. H ⁺	0.41	0.59	Ns
K _f vs Exch. Acidity	0.73	0.27	Ns
K _f vs Al ₂ O ₃	0.17	0.83	Ns
K _f vs Fe ₂ O ₃	0.15	0.85	Ns
1/n vs Clay	0.53	0.47	Ns
1/n vs pH	-0.88	0.12	Ns
1/n vs OM	-0.45	0.55	Ns
1/n vs Al ³⁺	1.00***	0.00001	***
1/n vs K _f	0.95*	0.05	*

Notes: *** = very strongly significant ($P \leq 0.001$); * = Significant at ($P \leq 0.05$); ns = non-Significant ($P > 0.05$); OM = Organic matter; Exch Al³⁺ = Exchangeable aluminium; Exch H⁺ = Exchangeable hydrogen; Exch Ac = Exchangeable acidity; Al₂O₃ = aluminium oxide; Fe₂O₃ = iron oxide; 1/n = sorption energy.

The K_f had a negative relation with the soil OM. When the soil OM increased, the K_f decreased. Similarly, Wolde and Haile (2015) stated, presence of OM decreased K_f of the soils due to the organic anions occupying the adsorption sites. Wehrauch and Opp (2018) also noted, organic acids chelate Fe/Al/Ca/Mg on oxide surfaces and hinder P sorption. For this particular study, a significantly ($P \leq 0.05$) positive relation between K_f and exch. Al^{3+} revealed, Al^{3+} was the main contributor for phosphate sorption in high P fixing soils (Alisols). The current finding is in concord with the findings of Hoseini and Taleshmikael (2013) described as insoluble Al phosphates are formed from the reaction of phosphate ions with Al^{3+} . A positive relationship between K_f and exch. H^+ indicates phosphate becomes less available at lower soils pH. Also, K_f had a positive relationship with Al_2O_3 and Fe_2O_3 as these oxides much contributed toward the increased K_f of the soils (Table 3). In addition, Al and Fe have a very high K_f in pure systems compared to their oxide forms since the oxides of Fe and Al are important cementing agents in soils which reduce the total surface area; therefore absorption sites (Reza, 2014).

5. CONCLUSION

There were clear variations in P sorption capacity and, therefore, External Phosphorus Requirement between soils of the study areas. The highest K_f (413 mg kg^{-1}) was observed for Alisols and the lowest K_f (280 mg kg^{-1}) was for Nitisols, even though, statistically no significant difference in the external phosphorus requirement. The actual quantity of P required by Nitisols, Luvisols, Alisols, and Andosols were 25, 30, 32, and 26 (mg kg^{-1}) respectively. The previously blanket P fertilizer recommendation in the country without considering differences among soil types is inadequate for optimal crops production. From the current study, 50 kg P ha^{-1} , 60 kg P ha^{-1} , 56 kg P ha^{-1} and 52 kg P ha^{-1} can be recommended for optimum crop production on Nitisols, Luvisols, Alisols, and Andosols respectively.

Declaration of Competing Interest

The authors declare no competing financial or personal interests that may appear and influence the work reported in this paper.

References

- Afsar, M. Z., Hoque, S., & Osman, K. (2012). A comparison of the Langmuir, Freundlich and Temkin equations to describe phosphate sorption characteristics of some representative soils of Bangladesh. *International Journal of Soil Science*, 7(3), 91. <https://doi.org/10.3923/ijss.2012.91.99>
- Beretta, A. N., Silbermann, A. V., Paladino, L., Torres, D., Bassahun, D., Musselli, R., & García-Lamohte, A. (2014). Soil texture analyses using a hydrometer: modification of the Bouyoucos method. *Ciencia e investigación agraria*, 41, 263-271. http://www.scielo.cl/scielo.php?script=sci_arttext&pid=S0718-16202014000200013&nrm=iso
- Buresh, R. J., Sánchez, P. A., & Calhoun, F. (1997). *Replenishing soil fertility in Africa* (Vol. 51). Soil Science Society of America <https://doi.org/10.2136/sssaspecpub51>
- Dari, B., Nair, V. D., Colee, J., Harris, W. G., & Mylavarapu, R. (2015). Estimation of phosphorus isotherm parameters: a simple and cost-effective procedure [Original Research]. *Frontiers in Environmental Science*, 3(70). <https://doi.org/10.3389/fenvs.2015.00070>
- Elias, E. (2016). *Soils of the Ethiopian Highlands: Geomorphology and Properties*. CASCAPE Project, ALTERA, Wageningen University and Research Centre (Wageningen UR). The Netherlands.
- Elias, E., Okoth, P. F., & Smaling, E. M. A. (2019). Explaining bread wheat (*Triticum aestivum*) yield differences by soil properties and fertilizer rates in the highlands of Ethiopia. *Geoderma*, 339, 126-133. <https://doi.org/10.1016/j.geoderma.2018.12.020>
- Fan, S.-S., Chang, F.-H., Hsueh, H.-T., & Ko, T.-H. (2016). Measurement of Total Free Iron in Soils by H_2S Chemisorption and Comparison with the Citrate Bicarbonate Dithionite Method. *Journal of Analytical Methods in Chemistry*, 2016, 7213542. <https://doi.org/10.1155/2016/7213542>
- Getie, A., Kiflu, A., & Meteke, G. (2021). Phosphorus Sorption Characteristics of Luvisols and Nitisols in North Ethiopian Soils. *Applied and Environmental Soil Science*, 2021, 8823852. <https://doi.org/10.1155/2021/8823852>
- Gonzalez-Rodriguez, S., & Fernandez-Marcos, M. L. (2018). Phosphate sorption and desorption by two contrasting volcanic soils of equatorial Africa. *PeerJ*, 6, e5820. <https://doi.org/10.7717/peerj.5820>
- Hanyabui, E., Apori, S. O., Frimpong, K. A., Atiah, K., Abindaw, T., Ali, M., Asiamah, J. Y., & Byalebeka, J. (2020). Phosphorus sorption in tropical soils. *AIMS Agriculture and Food*, 5(4), 599-616. <https://doi.org/10.3934/agrfood.2020.4.599>
- Hoseini, Y., & Taleshmikael, R. D. (2013). Comparison of phosphorus adsorption isotherms in soil and its relation to soil properties. *International Journal of Agriculture*, 3(1), 163.
- Jiang, X., Bol, R., Willbold, S., Vereecken, H., & Klumpp, E. (2015). Speciation and distribution of P associated with Fe and Al oxides in aggregate-sized fraction of an arable soil. *Biogeosciences*, 12(21), 6443-6452. <https://doi.org/10.5194/bg-12-6443-2015>
- Karlun, E., Mamo, T., Bekele, T., Gameda, S., & Kidanu, S. (2013). Ethiopian Soil Information System towards improved fertilizer recommendations in Ethiopia. *Nutrient indices for categorization of fertilizer blends from EthioSIS District soil inventory data discussion paper*.
- Kisinyo, P., Othieno, C., Gudu, S., Okalebo, J., Opala, P., Maghanga, J., Ng'etich, W., Agalo, J., Opile, R., & Kisinyo, J. (2015). Phosphorus sorption and lime requirements of maize growing acid soils of Kenya. <https://doi.org/10.5539/sar.v2n2p116>
- Lestari, Y., Maas, A., Purwanto, B. H., & Utami, S. N. H. (2016). The influence of lime and nitrogen fertilizer on soil

- acidity, growth and nitrogen uptake of corn in total reclaimed potential acid sulphate soil. *Journal of Agricultural Science*, 8(12), 197-205. <https://doi.org/10.5539/jas.v8n12p197>
- Maluf, H. J. G. M., Silva, C. A., Curi, N., Norton, L. D., & Rosa, S. D. (2018). Adsorption and availability of phosphorus in response to humic acid rates in soils limed with CaCO₃ or MgCO₃. *Ciência e Agrotecnologia*, 42, 7-20. <https://doi.org/10.1590/1413-70542018421014518>
- Melese, A., Gebrekidan, H., Yli-Halla, M., & Yitafaru, B. (2015). Phosphorus Status, Inorganic Phosphorus Forms, and Other Physicochemical Properties of Acid Soils of Farta District, Northwestern Highlands of Ethiopia. *Applied and Environmental Soil Science*, 2015, 748390. <https://doi.org/10.1155/2015/748390>
- Muindi, E. M., Mrema, J., Semu, E., Mtakwa, P., & Gachene, C. (2015). Effects of lime-aluminium-phosphate interactions on maize growth and yields in acid soils of the Kenya highlands. *American Journal of Agriculture and Forestry*, 3(6), 244-252. <https://doi.org/10.11648/j.ajaf.20150306.11>
- Reza, M. A. (2014). *Phosphate Sorption Behavior in Bajoa and Gopalpur Soil Series* [Thesis for B.Sc(Hon's), Khulna University]. <https://doi.org/10.13140/2.1.2529.1044>
- SAS Institute Inc. (2019). *SAS/SHARE® 9.4: User's Guide, Second Edition*. https://documentation.sas.com/doc/en/pgmsascdc/9.4_3.5/shrref/titlepage.htm
- Shirvani, M., Shariatmadari, H., & Kalbasi, M. (2005). Phosphorus Buffering Capacity Indices as Related to Soil Properties and Plant Uptake. *Journal of Plant Nutrition*, 28(3), 537-550. <https://doi.org/10.1081/PLN-200049235>
- Tamungang, N. E. B., David, M.-Z. A., Alakeh, M. N., & Adalbert, O. A. (2014). Phosphorus adsorption isotherms in relation to soil characteristics of some selected volcanic affected soils of Foubot in the West Region of Cameroon. *International Journal of Soil Science*, 11(2), 19-28. <https://doi.org/10.3923/ijss.2016.19.28>
- Thuy, P. T. P., Hoa, N. M., & Dick, W. A. (2020). Reducing Phosphorus Fertilizer Input in High Phosphorus Soils for Sustainable Agriculture in the Mekong Delta, Vietnam. *Agriculture*, 10(3), 87. <https://www.mdpi.com/2077-0472/10/3/87>
- Towett, E. K., Shepherd, K. D., & Cadisch, G. (2013). Quantification of total element concentrations in soils using total X-ray fluorescence spectroscopy (TXRF). *Science of The Total Environment*, 463-464, 374-388. <https://doi.org/10.1016/j.scitotenv.2013.05.068>
- Wang, B., Sun, J. S., Liu, H., & Ma, Y. B. (2017). The characteristics of phosphorus adsorption and desorption in gray desert soil of Xinjiang, China. *IOP Conference Series: Earth and Environmental Science*, 77, 012020. <https://doi.org/10.1088/1755-1315/77/1/012020>
- Weihrauch, C., & Opp, C. (2018). Ecologically relevant phosphorus pools in soils and their dynamics: The story so far. *Geoderma*, 325, 183-194. <https://doi.org/10.1016/j.geoderma.2018.02.047>
- Wolde, Z., & Haile, W. (2015). Phosphorus sorption isotherms and external phosphorus requirements of some soils of southern Ethiopia. *African Crop Science Journal*, 23(2), 89-99. <https://www.ajol.info/index.php/acsj/article/view/117723>